

doi: 10.13241/j.cnki.pmb.2017.19.018

糖尿病肾病患者红细胞免疫及抗氧化状态相关指标的变化研究 *

朱智全¹ 张壹言² 时新慧³ 杨文海⁴ 闫 华¹

(1 原军委后勤保障部第二门诊部 北京 100071; 2 解放军总医院第一附属医院肾内科 北京 100048;

3 解放军总医院第一附属医院检验科 北京 100048; 4 原军委训练管理部第三门诊部 北京 100857)

摘要 目的: 研究与分析糖尿病肾病患者红细胞免疫及抗氧化状态相关指标的变化情况。**方法:** 将 2015 年 1 月~2016 年 6 月期间于本院进行治疗的 66 例糖尿病肾病患者设为观察组, 同时将 66 名同龄健康人设为对照组。检测与比较两组同龄者的红细胞免疫及抗氧化状态相关指标, 同时比较不同糖尿病肾病分期患者的红细胞免疫及抗氧化状态相关指标水平。**结果:** 观察组的红细胞免疫及抗氧化状态相关指标明显差于对照组, 且不同糖尿病肾病分期患者的检测水平也呈现明显差异, P 均 < 0.05 , 检测水平之间均存在显著性差异。**结论:** 糖尿病肾病患者红细胞免疫及抗氧化状态相关指标明显较差, 糖尿病分期较高者的检测结果也相对较差, 因此应注意对糖尿病肾病患者进行上述方面的改善。

关键词: 糖尿病肾病; 红细胞免疫; 抗氧化状态相关指标

中图分类号: R587.2 文献标识码: A 文章编号: 1673-6273(2017)19-3675-03

Study on Changes of Erythrocyte Immune and Antioxidant Indexes of Patients with Diabetic Nephropathy*

ZHU Zhi-quan¹, ZHANG Yi-yang², SHI Xin-hui³, YANG Wen-hai⁴, YAN Hua¹

(1 Second Out-patient Department, Military Logistics Support Bureau, Beijing, 100071, China;

2 Department of Nephrology, the First Hospital Affiliated to General Hospital of PLA, Beijing, 100048, China;

3 Department of Laboratory, the First Hospital Affiliated to General Hospital of PLA, Beijing, 100048, China;

4 Third Out-patient Department, Military Training Administration Bureau, Beijing, 100857, China)

ABSTRACT Objective: To study and analyze the changes of erythrocyte immune and antioxidant indexes of patients with diabetic nephropathy. **Methods:** 66 patients with diabetic nephropathy in our hospital from Jan 2015 to June 2016 were chosen as observation group. 66 healthy persons of the same age were selected as the control group. Then the erythrocyte immune and antioxidant indexes of two groups with the same age were detected and compared. Also, the erythrocyte immune and antioxidant indexes levels of patients with different stages of diabetic nephropathy were compared. **Results:** The erythrocyte immune and antioxidant indexes of observation group were all obviously worse than those of control group. The detection levels of patients with different stages of diabetic nephropathy were all with significant differences too, all $P < 0.05$. **Conclusion:** The erythrocyte immune and antioxidant indexes of patients with diabetic nephropathy are obviously worse, and the detection results of patients with higher stages of diabetic nephropathy are relatively worse. Therefore, more attention should be paid to improve these aspects in patients with diabetic nephropathy.

Key words: Diabetic nephropathy; Erythrocyte immune; Antioxidant indexes

Chinese Library Classification(CLC): R587.2 **Document code:** A

Article ID: 1673-6273(2017)19-3675-03

前言

糖尿病肾病是糖尿病最重要的合并症之一, 近年来在我国发病率呈持续上升趋势, 已成为终末期肾脏疾病的第二大原因, 仅次于肾小球肾炎。由于糖尿病肾病代谢紊乱十分复杂, 如不加有效干预, 发展为终末期肾脏病后更加难以治愈。关于糖尿病肾病患者的研究显示^[1,2], 此类患者的免疫及血管受损状态均较为明显, 而机体的抗氧化能力的减弱可作为血管受损的重要因素, 对患者血液循环和新陈代谢的影响较大。同时, 机体免

疫评估中红细胞免疫指标可作为一项重要的代表性指标, 在糖尿病肾病患者中的具有较高研究意义^[3,4]。本文中我们就糖尿病肾病患者红细胞免疫及抗氧化状态相关指标的变化情况进行研究与分析, 现总结报道如下。

1 资料与方法

1.1 临床资料

选取 2015 年 1 月~2016 年 6 月期间于本院就诊的 66 例糖尿病肾病患者为研究对象, 设为观察组, 其中男性 37 例, 女

* 基金项目: 北京市科委 -- 首都医学发展科研基金项目(2007-3024)

作者简介: 朱智全(1971-), 本科, 研究方向: 临床检验指标在肝癌、肝硬化的检测价值等, E-mail: zhuzhiq6212@sina.com

(收稿日期: 2017-01-02 接受日期: 2017-01-26)

性 29 例;年龄为 36~74 岁,平均年龄为(54.5± 6.8)岁;病程为 1.8~18.5 年,平均病程为(9.9± 1.6)年;疾病分期:Ⅰ期者 12 例,Ⅱ期者 14 例,Ⅲ期者 14 例,Ⅳ期者 14 例,Ⅴ期者 12 例。本文对照组选取了 66 名同龄健康人,其中男性 36 名,女性 30 名;年龄为 36~75 岁,平均年龄为(54.2± 7.0)岁。两组的男女比例、年龄等均无显著性差异,具有可比性($P>0.05$)。

1.2 方法

所有研究对象于晨起空腹状态下采集 5.0 mL 的外周静脉血,离心后采集血清进行抗氧化状态相关指标的检测,检测项目为血清 TAC、CAT、LPO 及 MDA,检测方法为 ELISA 定量检测,另以郭峰法检测红细胞免疫指标,包括 ATER、FEER、FEIR 及 NTER。统计与比较两组同龄者的红细胞免疫及抗氧化状态相关指标,同时比较不同糖尿病肾病分期患者的红细胞免疫及

抗氧化状态相关指标水平。

1.3 统计学检验

本研究中的数据检验软件采用 SPSS20.0,计量资料的表示为 $\bar{x} \pm s$,计数资料的表示为率(%),上述两类数据分别以 t 检验和卡方检验分析, $P<0.05$ 表示比较结果间有显著性差异。

2 结果

2.1 两组同龄者的红细胞免疫及抗氧化状态相关指标检测结果比较

观察组的红细胞免疫指标 ATER、FEER 及 NTER 均低于对照组,FEIR 则高于对照组,抗氧化指标中的 TAC 及 CAT 均低于对照组,血清 LPO 及 MDA 则低于对照组, P 均 <0.05 ,具体比较见表 1。

表 1 两组同龄者的红细胞免疫检测结果比较

Table 1 Comparison of erythrocyte immune detection results of patients with the same age in the two groups

Detection indexes	Control group (n=66)	Observation group (n=66)
Erythrocyte immune index	ATER(%)	60.12± 5.66
	FEER(%)	66.56± 6.32
	FEIR (%)	22.10± 2.54
	NTER(%)	2.35± 0.24
Antioxidant indexes	TAC(U/ml)	13.23± 1.56
	CAT(nU/ml)	47.53± 5.12
	LPO(nmol/ml)	2.02± 0.30
	MDA(nmol/ml)	1.40± 0.21

Note: compared with control group, ° $P<0.05$.

2.2 不同糖尿病肾病分期患者的红细胞免疫及抗氧化状态相关指标检测结果比较

分期较高的糖尿病肾病患者的红细胞免疫指标 ATER、

FEER 及 NTER 均低于分期较高者,FEIR 则高于分期较高者,抗氧化指标中的 TAC 及 CAT 均低于分期较高者,血清 LPO 及 MDA 则低于分期较高者, P 均 <0.05 ,具体比较见表 2。

表 2 不同糖尿病肾病分期患者的红细胞免疫及抗氧化状态相关指标检测结果比较

Table 2 Comparison of erythrocyte immune and antioxidant indexes of patients with different stages of diabetic nephropathy

Detection indexes		Stage I	Stage II	Stage III	Stage IV	Stage V
Erythrocyte immune index	ATER(%)	58.68± 5.32	54.61± 5.10 ^o	51.02± 4.64 ^o	47.82± 4.46 ^o	43.20± 3.98 ^o
	FEER(%)	63.37± 6.12	60.18± 6.01 ^o	57.22± 5.46 ^o	54.32± 5.22 ^o	51.63± 4.80 ^o
	FEIR (%)	23.41± 2.65	26.45± 2.84 ^o	30.24± 3.11 ^o	33.56± 3.45 ^o	37.18± 3.74 ^o
	NTER(%)	2.18± 0.20	1.72± 0.16 ^o	1.20± 0.11 ^o	0.95± 0.08 ^o	0.67± 0.05 ^o
Antioxidant index	TAC(U/mL)	12.67± 1.48	10.38± 1.20 ^o	8.41± 0.95 ^o	6.62± 0.72 ^o	5.01± 0.54 ^o
	CAT(nU/mL)	44.28± 4.89	41.23± 4.66 ^o	38.10± 4.46 ^o	34.24± 4.10 ^o	30.05± 3.84 ^o
	LPO(nmol/mL)	2.21± 0.35	3.10± 0.45 ^o	4.13± 0.44 ^o	5.02± 0.50 ^o	6.11± 0.57 ^o
	MDA(nmol/mL)	1.54± 0.23	2.02± 0.32 ^o	3.01± 0.38 ^o	3.97± 0.40 ^o	4.85± 0.46 ^o

Note: compared with patients in stage I ~IV, ° $P<0.05$; compared with patients in stage I ~III, ° $P<0.05$; compared with patients in stage I ~II, ° $P<0.05$; compared with patients in stage I, ° $P<0.05$.

3 讨论

糖尿病肾病是糖尿病常见的糖尿病并发症,关于本病患者的诊断、检查及治疗方面的研究所占比例较高,而众多与之相关的研究显示,此类患者的血管状态及肾功能状态处于较差的状态^[5-8],而这些方面均受机体的氧化应激影响较大,而抗氧化

功能对于机体受损情况的修复等具有积极的临床意义,同时抗氧化功能状态与血管功能状态的关系极为密切,同时期还与机体的免疫状态有较大的关系,因此对于糖尿病肾病患者进行抗氧化能力状态的变化研究意义较高,可进一步了解机体的损伤程度及修复损伤的能力^[9-12],也可从一个侧面了解机体的血管功能状态。再者,红细胞免疫作为机体的一类有效防御机制,其

对于机体的免疫复合物清除具有极高的价值,临床对于重视程度不断提升的同时,与本类免疫相关的研究也持续增多,较多患者包括多类慢性病患者的此方面状态较差^[13-16],而关于糖尿病肾病患者红细胞免疫异常的研究也较多,但是关于其细致的研究十分不足,且研究差异的程度较大,故认为此方面的研究仍极为必要,以为糖尿病肾病患者免疫调节提供依据。

本文中我们就糖尿病肾病患者红细胞免疫及抗氧化状态相关指标的变化情况研究分析的结果显示,糖尿病肾病患者上述检测指标水平平均明显差于健康者,同时糖尿病肾病分期越高的患者其表达水平也相对越差,说明此类患者存在明显的免疫低下及抗氧化能力受损的情况,因此此类患者的血管功能状态相对较差,除与高血糖对血管造成的损伤有关外,机体抗氧化状态受损也是导致自由基清除较差的重要原因,而这进一步对血管造成损伤,因此血管功能较差^[17-20]。

综上所述,我们认为糖尿病肾病患者红细胞免疫及抗氧化状态相关指标明显较差,糖尿病分期较高者的检测结果也相对较差,因此应助于对糖尿病肾病患者进行上述方面的改善。

参考文献(References)

- [1] 陈敏,程丽霞.抗氧化应激治疗糖尿病肾病研究进展[J].青岛医药卫生, 2016, 48(4): 292-294
Chen Min, Cheng Li-xia. Research progress of oxidation stress treatment of diabetic nephropathy [J]. Qingdao Medical Journal, 2016, 48 (4): 292-294
- [2] 毛志敏,黄燕如,万毅刚,等.糖尿病肾病肾组织氧化应激的调控机制及中药的干预作用[J].中国中药杂志, 2014, 39(19): 3707-3712
Mao Zhi-min, Huang Yan-ru, Wan Yi-gang, et al. Regulative mechanisms of oxidative stress in kidney in diabetic nephropathy and interventional effects of Chinese herbal medicine[J]. China Journal of Chinese Materia Medica, 2014, 39(19): 3707-3712
- [3] 宋碧辉,谢红萍.氧化应激和 Nrf2 在糖尿病肾病发病机制中的研究进展[J].医学新知杂志, 2014, 14(5): 324-326
Song Bi-hui, Xie Hong-ping. Research progress of oxidative stress and Nrf2 in pathogenesis of diabetic nephropathy [J]. Journal of New Medicine, 2014, 14(5): 324-326
- [4] 郭福团,许雄伟,潘建峰,等.桑枝多糖对糖尿病肾病小鼠肾皮质氧化应激作用的影响[J].中国药理学通报, 2016, 32(8): 1148-1152
Guo Fu-tuan, Xu Xiong-wei, Pan Jian-feng, et al. The anti-oxidative effect of Ramulus mori polysaccharides on diabetic nephropathy mice [J]. Chinese Pharmacological Bulletin, 2016, 32(8): 1148-1152
- [5] 王国馨,唐英,王琛,等.糖肾宁 2 号方治疗 IV 期糖尿病肾病临床疗效及其对氧化应激指标的影响[J].中国中医药信息杂志, 2014, 14(5): 40-42, 49
Wang Guo-xin, Tang Ying, Wang Chen, et al. Clinical Efficacy of No.2 Tangshenning Decoction in Treating IV Stage Diabetic Nephropathy and Its Effect on Oxidative Stress Indicators[J]. Chinese Journal of Information on Traditional Chinese Medicine, 2014, 14(5): 40-42, 49
- [6] 王晓光,谭婷婷,任迪,等.补脾益肾活血法对糖尿病肾病患者氧化应激水平及炎症因子的影响[J].中国中西医结合肾病杂志, 2015, 15 (12): 1063-1067
Wang Xiao-guang, Tan Ting-ting, Ren Di, et al. Effect of Spleen-and-kidney-tonifying and Blood-circulating Therapy on the Oxidative Stress and Inflammatory Factors of Patients with Diabetic Nephropathy [J]. Chinese Journal of Integrated Traditional and Western Nephrology, 2015, 15(12): 1063-1067
- [7] 杨林燕,李英.抗氧化应激对糖尿病肾病肾脏损害的保护作用[J].中国中西医结合肾病杂志, 2016, 17(6): 554-556
Yang Lin-yan, Li Ying. Protective function of antioxidant stress on renal damage in diabetic nephropathy [J]. Chinese Journal of Integrated Traditional and Western Nephrology, 2016, 17(6): 554-556
- [8] 田华,胡灵卫,张玉霞,等.姜黄素对糖尿病肾病大鼠肾脏的抗氧化应激作用及保护机制[J].中国临床药理学杂志, 2013, 29(6): 457-459
Tian Hua, Hu Ling-wei, Zhang Yu-xia, et al. Antioxidative effect of curcumin on diabetic nephropathy rats and its protective mechanism [J]. The Chinese Journal of Clinical Pharmacology, 2013, 29 (6): 457-459
- [9] 寿成珉,周旦阳,陈惠萍,等.糖尿病肾病血清抗氧化能力及炎症指标的变化[J].中华全科医学, 2014, 12(2): 245-247
Shou Cheng-min, Zhou Dan-yang, Chen Hui-ping, et al. Change of serum oxidation resistance and inflammatory markers of patients with diabetic nephropathy [J]. Chinese Journal of General Practice, 2014, 12(2): 245-247
- [10] 张红霞.京尼平对糖尿病肾病小鼠肾脏的保护作用[J].医药导报, 2015, 34(1): 26-30
Zhang Hong-xia. Protective Effects of Genipin on Diabetic Nephropathy in Mice[J]. Herald of Medicine, 2015, 34(1): 26-30
- [11] Perlman AS, Chevalier JM, Wilkinson P, et al. Serum Inflammatory and Immune Mediators Are Elevated in Early Stage Diabetic Nephropathy[J]. Ann Clin Lab Sci, 2015, 45(3): 256-263
- [12] 刘婧,扈会整,张勇,等.糖尿病肾病患者血浆 NGAL 和血清 CysC 水平改变及其早期诊断价值[J].现代生物医学进展, 2015, 15(30): 5945-5947, 5900
Liu Jing, Hu Hui-zheng, Zhang Yong, et al. Changes of Serum NGAL and CysC Levels in Patients with Diabetic Nephropathy and Their Early Diagnostic Value[J]. Progress in Modern Biomedicine, 2015, 15 (30): 5945-5947, 5900
- [13] 杨楠楠,刚晓坤,刘青,等.糖尿病肾病与氧化应激[J].中国老年学杂志, 2013, 33(5): 1237-1239
Yang Nan-nan, Gang Xiao-kun, Liu Qing, et al. Diabetic nephropathy and oxidative stress [J]. Chinese Journal of Gerontology, 2013, 33(5): 1237-1239
- [14] 张玉霞,刘国玲,王家勤,等.雷公藤多苷对糖尿病肾病大鼠的抗氧化应激作用[J].中国药理学与毒理学杂志, 2014, 14(3): 358-361
Zhang Yu-xia, Liu Guo-ling, Wang Jia-qin, et al. Antioxidative effect of Tripterygium wilfordii polyglycosides on diabetic rats [J]. Chinese Journal of Pharmacology and Toxicology, 2014, 14(3): 358-361
- [15] 张体华,王明.糖尿病肾病患者炎症水平、免疫功能及与肾脏病变的关系[J].中国免疫学杂志, 2016, 32(10): 1524-1526, 1531
Zhang Ti-hua, Wang Ming. Relationship of micro-inflammation, cellular and humoral immune function and kidney disease of patients with diabetic nephropathy [J]. Chinese Journal of Immunology, 2016, 32(10): 1524-1526, 1531
- [16] 张洋,马坤岭,刘晶,等.慢性炎症反应对糖尿病肾病的促进作用[J].中华肾脏病杂志, 2013, 29(9): 681-686
Zhang Yang, Ma Kun-ling, Liu Jing, et al. Chronic inflammatory response promotes diabetic nephropathy [J]. Chinese Journal of Nephrology, 2013, 29(9): 681-686

- [4] Xu H, Li S. Pharmacological effects of Bailing capsule and its application in lung disease research [J]. Zhongguo Zhong Yao Za Zhi, 2010, 35(20): 2777-2781
- [5] Shahebrahimi K, Jalilian N, Bazgir N, et al. Comparison clinical and metabolic effects of metformin and pioglitazone in polycystic ovary syndrome[J]. Indian J Endocrinol Metab, 2016, 20(6): 805-809
- [6] Adamska A, Łebkowska A, Krentowska A, et al. Relationship between serum gonadotropin concentrations and thyroid volume in women with polycystic ovary syndrome [J]. Pol Arch Med Wewn, 2016, 126(11): 891-894
- [7] Luz R, Barros J, Aguiar A, et al. Short and Long-Term Efficacy of Laparoscopic Ovarian Diathermy in Women with Polycystic Ovary Syndrome[J]. Acta Med Port, 2016, 29(7-8): 441-448
- [8] Cassar S, Misso ML, Hopkins WG, et al. Insulin resistance in polycystic ovary syndrome: a systematic review and meta-analysis of euglycaemic-hyperinsulinaemic clamp studies [J]. Hum Reprod, 2016, 31(11): 2619-2631
- [9] Park CH, Chun S. Association between serum gonadotropin level and insulin resistance-related parameters in Korean women with polycystic ovary syndrome[J]. Obstet Gynecol Sci, 2016, 59(6): 498-505
- [10] Dube R. Does endothelial dysfunction correlate with endocrinal abnormalities in patients with polycystic ovary syndrome?[J]. Avicenna J Med, 2016, 6(4): 91-102
- [11] Banaszewska B, Wrotyńska-Barczyńska J, Spaczynski RZ, et al. Effects of Resveratrol on Polycystic Ovary Syndrome: A Double-blind, Randomized, Placebo-controlled Trial[J]. J Clin Endocrinol Metab, 2016, 101(11): 4322-4328
- [12] Song J, Li YH, Yang XD. Effect of combined therapy with bailing capsule and benazepril on urinary albumin excretion rate and C-reactive protein in patients with early diabetic nephropathy [J]. Zhongguo Zhong Xi Yi Jie He Za Zhi, 2009, 29(9): 791-793
- [13] Wang W, Zhang XN, Yin H, et al. Effects of Bailing capsules for renal transplant recipients: a retrospective clinical study[J]. Chin Med J (Engl), 2013, 126(10): 1895-1899
- [14] Wei LN, Huang R, Li LL, et al. Reduced and delayed expression of GDF9 and BMP15 in ovarian tissues from women with polycystic ovary syndrome[J]. J Assist Reprod Genet, 2014, 31(11): 1483-1190
- [15] Persani L, Rossetti R, Di Pasquale E, et al. The fundamental role of bone morphogenetic protein 15 in ovarian function and its involvement in female fertility disorders [J]. Hum Reprod Update, 2014, 20(6): 869-883
- [16] Berberoglu Z, Aktas A, Fidan Y, et al. Association of plasma GDF-9 or GDF-15 levels with bone parameters in polycystic ovary syndrome [J]. J Bone Miner Metab, 2015, 33(1): 101-108
- [17] Wei LN, Fang C, Huang R, et al. Change and significance of growth differentiation factor 9 and bone morphogenetic protein expression during oocyte maturation in polycystic ovary syndrome patients with ovarian stimulation [J]. Zhonghua Fu Chan Ke Za Zhi, 2012, 47(11): 818-822
- [18] Luo L, Wang Q, Chen M, et al. IGF-1 and IGFBP-1 in peripheral blood and decidua of early miscarriages with euploid embryos: comparison between women with and without PCOS [J]. Gynecol Endocrinol, 2016, 32(7): 538-542
- [19] Brazert M, Pawelczyk LA. Insulin-like growth factor-1 isoforms in human ovary. Preliminary report on the expression of the IGF-1 gene in PCOS patients and healthy controls[J]. Ginekol Pol, 2015, 86(12): 890-895
- [20] Shahnazi V, Zaree M, Nouri M, et al. Influence of ω -3 fatty acid eicosapentaenoic acid on IGF-1 and COX-2 gene expression in granulosa cells of PCOS women[J]. Iran J Reprod Med, 2015, 13(2): 71-78

(上接第 3677 页)

- Zhang Yang, Ma Kun-ling, Liu Jing, et al. Inflammatory stress exacerbates the progression of diabetic nephropathy[J]. Chinese Journal of Nephrology, 2013, 29(9): 681-686
- [17] 田勇, 王俊宏. 前列地尔对老年糖尿病肾病患者氧化应激及免疫功能的影响[J]. 中国老年学杂志, 2015, 15(9): 2419-2421
- Tian Yong, Wang Jun-hong. Influence of Alprostadil on oxidative stress and immune function of diabetic nephropathy in elderly patients[J]. Chinese Journal of Gerontology, 2015, 15(9): 2419-2421
- [18] 戴爱明, 李翠兰. 前列地尔联合阿卡波糖治疗 2 型糖尿病肾病的疗效及对氧化应激水平的影响 [J]. 医学临床研究, 2015, 15(1): 113-115
- Dai Ai-ming, Li Cui-lan. Curative Efficacy of Prostaglandin Plus Acarbose and its Effects on Oxidative Stress in Patients with Type 2 Diabetic Nephropathy [J]. Journal of Clinical Research, 2015, 15(1): 113-115
- [19] Korabecna M, Pazourkova E, Horinek A, et al. Methylation status of immune response genes promoters in cell-free DNA differs in hemodialyzed patients with diabetic nephropathy according to the intensity of anemia therapy[J]. Blood Purif, 2013, 36(3-4): 280-286
- [20] 雍军光, 阮萍, 冯博华, 等. 糖尿病肾病患者红细胞形态功能变化与病程发展的关联性研究[J]. 暨南大学学报(自然科学与医学版), 2013, 13(6): 610-614
- Yong Jun-guang, Yuan Ping, Feng Bo-hua, et al. Study on the relevance of changes in red blood cell morphology and function of patients with diabetic nephropathy to the development of diabetes [J]. Journal of Jinan University (Natural Science & Medicine Edition), 2013, 13(6): 610-614