

doi: 10.13241/j.cnki.pmb.2018.09.034

快速康复外科理念对结直肠癌根治术患者疗效及机体应激反应的影响 *

甘 涛¹ 夏 涛^{1△} 李 威¹ 杨 颢¹ 莫松江²

(1 广西科技大学附属柳州市人民医院胃肠外科 广西 柳州 545006;

2 广西柳州市鹿寨县人民医院普通外科 广西 柳州 545600)

摘要 目的: 探讨快速康复外科理念应用于结直肠癌根治术患者的疗效及其对机体应激反应的影响。**方法:** 选取 2012 年 1 月-2013 年 6 月我院收治的 180 例结直肠癌患者为研究对象,采用随机数字表法分为对照组与观察组,每组各 90 例。两组患者均进行结直肠癌根治术治疗,对照组患者围手术期采用传统处理措施,观察组患者围手术期采用快速康复处理措施,对比两组患者术中出血量、手术时间、术后住院时间、住院费用、术后排便时间、首次排气时间,同时比较两组患者手术当日、术后 1 d、5 d 血清白细胞介素-6(IL-6)、C 反应蛋白(CRP)及血清淀粉样蛋白 A(SAA)的水平,并观察两组患者并发症发生情况。**结果:** 与对照组相比,观察组患者的术后住院时间、术后排便时间及首次排气时间均缩短,住院费用降低,差异有统计学意义($P<0.05$),两组患者术中出血量、手术时间对比差异无统计学意义($P>0.05$)。术前,两组患者的血清 IL-6、CRP、SAA 水平对比差异无统计学意义($P>0.05$),术后 1 d、5 d,两组患者血清 IL-6、CRP、SAA 水平均高于术前,术后 5 d 血清 IL-6、CRP、SAA 水平低于术后 1 d,差异有统计学意义($P<0.05$),观察组患者术后 1 d、5 d 血清 IL-6、CRP、SAA 水平均低于对照组,差异有统计学意义($P<0.05$)。观察组并发症总发生率为 6.67%,与对照组的 7.78%比较差异无统计学意义($P>0.05$)。**结论:** 快速康复外科理念应用于结直肠癌根治术患者能够有效加快患者术后康复,降低术后应激反应,值得临床推广。

关键词: 结直肠癌根治术;快速康复外科理念;疗效;白细胞介素-6;C 反应蛋白;血清淀粉样蛋白 A

中图分类号: R735.3 **文献标识码:** A **文章编号:** 1673-6273(2018)09-1758-04

Application of Concept of Rapid Rehabilitation Surgery in Patients with Radical Resection of Colorectal Cancer*

GAN Tao¹, XIA Tao^{1△}, LI Wei¹, YANG Hao¹, MO Song-jiang²

(1 Department of Gastrointestinal Surgery, Liuzhou People's Hospital Affiliated to Guangxi University of Science and Technology, Liuzhou, Guangxi, 545006, China;

2 Department of General Surgery, Luzhai County People's Hospital of Liuzhou, Liuzhou, Guangxi, 545600, China)

ABSTRACT Objective: To explore the Effect of rapid rehabilitation surgery concept on the radical resection of colorectal cancer, and to analyse the stress response on the body. **Methods:** A total of 180 cases of colorectal cancer, who were treated in Liuzhou People's Hospital Affiliated to Guangxi University of Science and Technology from January 2012 to June 2013, were randomly divided into control group and observation group, 90 cases in each group. The two groups were treated with radical resection of colorectal cancer. The traditional treatment measures were adopted in the perioperative period of the control group; the measures of rapid rehabilitation surgery concept was used for the observation group during the perioperative period. The amount of bleeding, operation time, postoperative hospital stay, the cost of hospitalization, the time of postoperative defecation and the first exhaust time were compared between the two groups. At the same time, the levels of serum interleukin-6 (IL-6), C reactive protein (CRP) and serum amyloid A (SAA) in two groups of patients at operative time, 1 d and 5 d after operation were compared between the two groups. The complications in two groups were observed. **Results:** Compared with the control group, the time of postoperative hospital stay, the time of postoperative defecation and the first exhaust time were shortened, and the cost of hospitalization was reduced in the observation group, the differences were statistically significant($P<0.05$). There were no significant differences in the amount of bleeding and operation time between the two groups ($P>0.05$). Before operation, there were no significant differences in serum IL-6, CRP and SAA levels between the two groups ($P>0.05$); 1 d and 5 d after operation, the levels of serum IL-6, CRP and SAA in the two groups were all higher than those before operation, the levels of IL-6, CRP and SAA 5d after operation were lower than those 1 d after operation, the differences were statistically significant ($P<0.05$). The levels of serum IL-6, CRP and SAA in the observation group 1 d and 5 d after operation were lower than those in the control group, the difference was statistically significant ($P<0.05$). The total incidence of complications in the observation group was 6.67%, which has no signifi-

* 基金项目:广西壮族自治区卫生厅自筹经费科研课题(Z2012576)

作者简介:甘涛(1978-),男,本科,副主任医师,从事胃肠道肿瘤诊治方面的研究, E-mail:kwghdy@163.com

△ 通讯作者:夏涛(1962-),男,本科,副主任医师,从事胃肠道肿瘤诊治方面的研究, E-mail:hggyye@163.com

(收稿日期:2017-12-17 接受日期:2017-12-31)

cant difference compared with the total incidence of complications (7.78%) in the control group ($P>0.05$). **Conclusion:** The rapid rehabilitation surgery concept in the patients with radical resection of colorectal cancer can effectively accelerate the recovery of patients after operation and reduce postoperative stress response, which is worthy of clinical promotion.

Key words: Radical resection of colorectal cancer; Rapid rehabilitation surgery concept; Curative effect; Interleukin -6; C reactive protein; Serum amyloid A

Chinese Library Classification(CLC): R735.3 **Document code:** A

Article ID: 1673-6273(2018)09-1758-04

前言

结直肠癌是临床常见的消化道恶性肿瘤之一,具有较高的发病率及病死率,对人类的健康造成严重威胁^[1,2]。腹腔镜下结直肠癌根治术是目前临床上治疗结直肠癌的主要方法,该手术的围手术期肠道准备对手术效果影响较大^[3,4]。传统的围手术期处理包括清洁灌肠、术前肠道准备、术后留置胃肠减压、保留腹腔引流管和尿管及应用度冷丁镇痛等,以上围手术期处理极易增加术后疼痛感,同时会导致机体内环境紊乱,因此探寻更有效的围手术期肠道处理方法是临床工作者关注的重点^[5,6]。快速康复外科理念是近年来兴起的围手术期处理方法,其具体内容包括良好的麻醉、术前宣教、止痛、术后早期康复治疗以及围手术期各种应激反应治疗等,可以更好的促进患者术后胃肠功能的恢复,减轻患者术后疼痛,缩短患者恢复时间^[7,8]。本研究旨在探讨快速康复外科理念应用于结直肠癌根治术患者的疗效以及其对机体应激反应的影响。

1 资料与方法

1.1 一般资料

选取 2012 年 1 月 -2013 年 6 月我院收治的 180 例结直肠癌患者为研究对象,纳入标准^[9]:(1)术前确诊为结直肠癌;(2)术前经影像学检查或体检无其他脏器转移;(3)可行结直肠癌根治术;(4)患者及其家属知情本研究并签署知情同意书。排除标准:(1)伴有免疫系统及内分泌系统疾病的患者;(2)伴有肝肾功能及凝血功能障碍者;(3)伴有其他炎症性疾病者;(4)在术前接受过新辅助化疗治疗者。采用随机数字表法分为观察组($n=90$)与对照组($n=90$)。其中观察组男 45 例,女 45 例,年龄 43-69 岁,平均年龄(56.81 ± 4.34)岁;肿瘤位置:结肠癌 37 例,直肠癌 53 例;TMN 分期:I 期 21 例,II 期 47 例,III 期 22 例。对照组男 49 例,女 41 例,年龄 45-70 岁,平均年龄(58.93 ± 6.33)岁;肿瘤位置:结肠癌 40 例,直肠癌 50 例;TMN 分期:I 期 25 例,II 期 48 例,III 期 17 例。两组患者一般资料对比无差异($P>0.05$),提示分组均衡,组间具有可比性,本次研究通过医院伦理委员会批准。

1.2 围手术期处理方法

观察组患者围手术期给予快速康复外科简易肠道处理:患者在入院后正常饮食,在术前根据快速康复注意事项、处理流程及结直肠癌的相关知识进行健康宣教。术前禁食、禁饮 4 h,术前 1 d 口服复方聚乙二醇电解质散 2000 mL 经口全肠道清洁灌肠,然后口服肠内营养乳剂,采取连续硬膜外联合气管插管下全身麻醉,术后采用静脉自控式镇痛,手术次日即可流质饮食,术后 48 h 拔除导尿管。对照组患者围手术期采用传统处

理方法:开始手术前对患者进行常规宣教,在进行手术前 5 d 患者开始使用半流质食物,手术前 3 d 开始饮用流质食物,同时口服缓泻剂如硫酸镁,50 mL/d,每天晚上应用 2000 mL 的生理盐水经肛门进行机械性灌肠;在手术前 2-3 d 服用肠道抗生素:甲硝唑 0.4 g,3 次/d,链霉素 0.5 d,1 次/d;术前 1 d 禁食,术前当晚使用生理盐水经肛门行清洁灌肠,至排出大便为清水样,麻醉方式采用气管插管下全身麻醉,常规放置腹腔引流管,术后镇痛给予肌肉注射哌替啶或曲马多 50 mg,根据患者的疼痛程度按需止痛,肠道排气后开始饮水进食,术后 3-4 d 拔除导尿管。

1.3 观察指标及检测方法

对比两组患者术中出血量、手术时间、术后住院时间、住院费用、术后排便时间、首次排气时间,所有患者在手术当日、术后 1 d、5 d 采取外周空腹静脉血 5 mL,置于预冷试管内,应用离心机(美国 Beckman 公司)以 3000 r/min 的转速离心 10 min,将血清与血浆分离,将离心管上层的血清收集,置于 -70℃ 冰箱内保存待检。采用酶联免疫吸附法检测患者血清白细胞介素 -6 (Interleukin -6, IL-6)、C 反应蛋白(C reactive protein, CRP)及血清淀粉样蛋白 A(Serum amyloid A protein, SAA)的水平,试剂盒由上海盈公生物技术有限公司提供,操作按照试剂盒说明书进行。观察两组患者并发症发生情况。

1.4 统计学方法

数据分析应用 SPSS23.0 统计学软件,计数资料以%表示,采用 χ^2 检验;计量资料以($\bar{x}\pm s$)表示,采用 t 检验,检验标准设置为 $\alpha=0.05$ 。

2 结果

2.1 两组患者手术指标及术后恢复情况比较

与对照组相比,观察组患者的术后住院时间、术后排便时间及首次排气时间均缩短,住院费用降低($P<0.05$),两组患者术中出血量、手术时间对比差异无差异($P>0.05$),见表 1。

2.2 两组患者血清 IL-6、CRP、SAA 水平比较

术前,两组患者的血清 IL-6、CRP、SAA 水平对比差异无统计学意义($P>0.05$),术后 1 d、5 d, 两组患者血清 IL-6、CRP、SAA 水平均高于术前,术后 5 d 血清 IL-6、CRP、SAA 水平低于术后 1 d ($P<0.05$), 观察组患者术后 1 d、5 d 血清 IL-6、CRP、SAA 水平均低于对照组($P<0.05$),见表 2。

2.3 两组患者并发症发生情况比较

观察组并发症总发生率为 6.67%(6/90), 与对照组的 7.78%(7/90)比较差异无统计学意义($P>0.05$),见表 3。

表 1 两组患者手术指标及术后恢复情况比较($\bar{x} \pm s$)

Table 1 Comparison of surgical indexes and postoperative recovery in the two groups($\bar{x} \pm s$)

Groups	n	Amount of bleeding(mL)	Operation time (min)	Postoperative hospital stay(d)	Cost of hospitalization (million)	Time of postoperative defecation(h)	The first exhaust time(h)
Observation group	90	65.73± 15.14	115.79± 20.67	11.19± 1.92	2.63± 0.32	72.18± 21.33	62.15± 15.67
Control group	90	68.32± 19.46	120.53± 23.83	14.21± 3.25	3.76± 0.57	115.74± 28.75	85.54± 20.18
t		0.649	0.401	5.064	2.217	5.077	4.189
P		0.517	0.697	0.027	0.042	0.031	0.035

表 2 两组患者血清 IL-6、CRP、SAA 水平比较($\bar{x} \pm s$)

Table 2 Comparison of IL-6, CRP and SAA levels between the two groups($\bar{x} \pm s$)

Groups	n	IL-6(pg/L)			CRP(mg/L)			SAA(mg/L)		
		before operation	1 d after operation	5 d after operation	before operation	1 d after operation	5 d after operation	before operation	1 d after operation	5 d after operation
Observation group	90	3.78± 1.02	130.01± 7.54*	11.47± 3.32* [#]	2.34± 0.67	42.80± 7.39*	18.15± 5.38* [#]	3.53± 1.11	134.06± 16.89*	30.54± 7.32* [#]
		4.09± 1.27	154.19± 14.16*	21.21± 4.65* [#]	2.31± 0.54	79.29± 13.96*	40.64± 12.19* [#]	3.50± 1.05	182.38± 41.07*	58.59± 11.56* [#]
Control group	90	0.763	7.980	7.966	0.492	11.364	6.212	0.316	5.125	7.602
		0.419	0.027	0.029	0.673	0.007	0.019	0.837	0.038	0.024

Note: Compared with before operation, *P<0.05. Compared with 1d after operation, [#]P<0.05.

表 3 两组患者并发症发生情况比较[n(%)]

Table 3 Comparison of complications of the two groups [n(%)]

Groups	n	Pulmonary infection	Urinary tract infection	Postoperative ileus	Anastomotic leak	Total incidence
Observation group	90	2(2.22)	0(0.00)	1(1.11)	3(3.33)	6(6.67)
Control group	90	3(3.33)	2(2.22)	0(0.00)	2(2.22)	7(7.78)
χ^2						2.292
P						0.164

3 讨论

结直肠癌是胃肠道中常见的恶性肿瘤,世界流行病学统计报道显示,在西欧、新西兰、澳大利亚、北美等地区的结肠癌发病率较高,居内脏肿瘤前二位^[10-12]。根据各地区的资料显示,随着近年来人们生活水平的改善,饮食结构改变,结直肠癌的发病率逐年上升,且青年人在其中所占的比例也在逐年升高^[13-15]。目前临床上治疗结直肠癌的方法主要是手术切除,由于大部分结直肠癌患者在接受手术治疗前存在一定程度的免疫功能减退以及营养不良现象,所以结直肠癌根治术围手术期处理方法的选择对患者的后期治疗及恢复至关重要^[16-18]。虽然目前已有较多的肠道准备方法,并且比以往肠道准备方法具有较大的改进,能够满足临床手术的基本需求,但仍效果欠佳,因此探寻更有效的肠道准备方法对促进结直肠癌根治术患者术后恢复具有重要意义。

传统的肠道准备方法中需要在术前限制饮食,致使患者术前营养状态较差及水电解质内环境紊乱,使患者对手术的耐受性降低,增加了术后发生并发症的几率,影响手术效果^[19-21]。近年来,随着胃肠外科设备、技术的进步和快速康复外科新理念的发展,结直肠癌患者得到了良好治疗。为加快患者术后恢复、

减轻患者负担、争取术后尽快得到后续治疗,临床工作者将快速康复外科新理念运用于结直肠癌患者术前肠道准备中^[22]。快速康复外科理念产生的主要基础是对手术应激反应的深入认识,其能有效减轻患者术后的应激反应,进而使加快患者术后康复^[23]。本次研究结果表明,观察组术后住院时间、术后排便时间及首次排气时间均较对照组缩短,住院费用较对照组降低(P<0.05),说明应用快速康复外科理念对结直肠癌根治术患者进行围手术期处理可有效缩短患者术后恢复时间,减轻患者经济压力。分析其主要原因是观察组患者在术前正常饮食,营养状态未下降,同时术前未服用抗生素,降低了术后发生假膜性小肠结肠炎、腹泻等疾病的风险;术前一天患者口服复方聚乙二醇电解质散行进行全肠道清洁灌肠,既不会对肠道微生物造成明显的影响,也不会扰乱正常的肠道菌群,此外应用聚乙二醇进行肠道准备,不会使结肠黏膜形态学发生变化,肠道清洁度优于传统肠道准备,避免了水电解质内环境紊乱^[24,25];术前口服肠内营养乳剂,在满足患者营养需求的同时,还可刺激患者胃肠道,促进胃肠蠕动及黏膜生长,加快患者术后胃肠道功能的恢复,因此观察组患者术后康复时间较对照组缩短。应激反应是机体在受到各种内外环境因素刺激时所出现的反应,IL-6、CRP 及 SAA 是三种特征性急性反应介质,其中 IL-6 是一

种促炎因子,也是机体在应激反应状态下介导免疫调节及炎症损伤的重要细胞因子之一,能够较早的反映机体损伤程度^[26,27]; CRP 是机体受到组织损伤时产生的急性蛋白,反映急性期炎症反应的重要指标^[28];SAA 是一种比 CRP 更为敏感的炎性指标,其在血清中的水平可反应组织损伤程度^[29]。本研究在对以上三种指标水平对比分析中发现,观察组患者术后 1 d、5 d 的 IL-6、CRP、SAA 水平均低于对照组($P<0.05$),说明观察组患者术后应激反应低于对照组,分析其原因主要是观察组患者麻醉方式采用全身麻醉联合连续硬膜外麻醉,且术后采用静脉自控镇痛,可有效减少皮质醇、儿茶酚胺的释放,使应激反应减轻^[30]。此外,观察组患者在术后早期经口饮食,可有效促进肠道蠕动,使肠道胃肠激素的分泌增加,维护了肠粘膜的功能,避免了肠道菌群移位的发生,使应激反应减轻。本研究对两组并发症分析中显示,两组并发症发生率无差异($P>0.05$),说明快速康复外科理念围手术期处理方法不会增加并发症的发生率,具有较高的安全性。

综上所述,快速康复外科理念应用于结直肠癌根治术患者可有效缩短患者术后康复时间、住院时间,减轻术后应激反应,减少患者住院费用,既能促进患者术后康复,又能减轻患者经济压力,同时又具有较高的安全性,值得临床推广。

参考文献(References)

- [1] Zhang J, Jiang Y, Zhu J, et al. Overexpression of long non-coding RNA colon cancer-associated transcript 2 is associated with advanced tumor progression and poor prognosis in patients with colorectal cancer[J]. *Oncol Lett*, 2017, 14(6): 6907-6914
- [2] Tian S, Lei HB, Liu YL, et al. The association between metformin use and colorectal cancer survival among patients with diabetes mellitus: An updated meta-analysis [J]. *Chronic Dis Transl Med*, 2017, 3(3): 169-175
- [3] Morończyk DA, Krasnodebski IW. Implementation of the fast track surgery in patients undergoing the colonic resection: own experience [J]. *Pol Przegl Chir*, 2011, 83(9): 482-487
- [4] Fugang W, Zhaopeng Y, Meng Z, et al. Long-term outcomes of laparoscopy vs. open surgery for colorectal cancer in elderly patients: A meta-analysis[J]. *Mol Clin Oncol*, 2017, 7(5): 771-776
- [5] Cui JH, Jiang WW, Liao YJ, et al. Effects of oxycodone on immune function in patients undergoing radical resection of rectal cancer under general anesthesia[J]. *Medicine (Baltimore)*, 2017, 96(31): e7519
- [6] Backes Y, de Vos Tot Nederveen Cappel WH, van Bergeijk J, et al. Risk for Incomplete Resection after Macroscopic Radical Endoscopic Resection of T1 Colorectal Cancer: A Multicenter Cohort Study[J]. *Am J Gastroenterol*, 2017, 112(5): 785-796
- [7] 谢正勇,程黎阳,张玉新,等.快速康复外科理念结合腹腔镜用于治疗结直肠疾病的效果[J]. *广东医学*, 2015, 36(18): 2805-2806
Xie Zheng-yong, Cheng Li-yang, Zhang Yu-xin, et al. Efficacy of rapid rehabilitation surgery combined with laparoscopy in the treatment of colorectal diseases[J]. *Guangdong Medical Journal*, 2015, 36(18): 2805-2806
- [8] 曾晓峰,牛维益.快速康复外科理念在治疗结直肠癌围术期的应用研究[J]. *重庆医学*, 2017, 46(15): 2051-2053
Zeng Xiao-feng, Niu Wei-yi. Clinical study on the concept of rapid rehabilitation surgery in perioperative period of colorectal cancer[J]. *Chongqing Medicine*, 2017, 46(15): 2051-2053
- [9] 中华医学会外科学分会腹腔镜与内镜外科学组,中国抗癌协会大肠癌专业委员会腹腔镜外科学组. 腹腔镜结直肠癌根治手术操作指南(2008 版)[J]. *中华胃肠外科杂志*, 2009, 12(3): 310-312
Laparoscopic and endoscopic surgery department of the Chinese Medical Association, China Cancer Association, colorectal cancer (Specialized Committee) laparoscopic surgery group. Laparoscopic radical resection of colorectal cancer (2008 Edition)[J]. *Chinese Journal of Gastrointestinal Surgery*, 2009, 12(3): 310-312
- [10] Spychalski M, Skulimowski A, Dziki A, et al. Colorectal endoscopic submucosal dissection (ESD) in the West - when can satisfactory results be obtained? A single-operator learning curve analysis[J]. *Scand J Gastroenterol*, 2017, 52(12): 1442-1452
- [11] Rho YS, Gilabert M, Polom K, et al. Comparing Clinical Characteristics and Outcomes of Young-onset and Late-onset Colorectal Cancer: An International Collaborative Study [J]. *Clin Colorectal Cancer*, 2017, 16(4): 334-342
- [12] Leshno A, Moshkowitz M, David M, et al. Prevalence of colorectal neoplasms in young, average risk individuals: A turning tide between East and West[J]. *World J Gastroenterol*, 2016, 22(32): 7365-7372
- [13] Boyce S, Nassar N, Lee CY, et al. Young-onset colorectal cancer in New South Wales: a population-based study [J]. *Med J Aust*, 2016, 205(10): 465-470
- [14] Kato T, Alonso S, Muto Y, et al. Clinical characteristics of synchronous colorectal cancers in Japan [J]. *World J Surg Oncol*, 2016, 14(1): 272
- [15] Douaiher J, Ravipati A, Grams B, et al. Colorectal cancer-global burden, trends, and geographical variations [J]. *J Surg Oncol*, 2017, 115(5): 619-630
- [16] Chen WK, Ren L, Wei Y, et al. General anesthesia combined with epidural anesthesia ameliorates the effect of fast-track surgery by mitigating immunosuppression and facilitating intestinal functional recovery in colon cancer patients [J]. *Int J Colorectal Dis*, 2015, 30(4): 475-481
- [17] 夏秀娟,李粉婷,刘佳,等.转移性结直肠癌抗血管生成靶向治疗的研究进展[J]. *现代生物医学进展*, 2016, 16(1): 187-191
Xia Xiu-juan, Li Fen-ting, Liu Jia, et al. Anti-angiogenesis Targeted Therapy for Metastatic Colorectal Cancer: Current Situation and Future Perspectives [J]. *Progress in Modern Biomedicine*, 2016, 16(1): 187-191
- [18] Li L, Jin J, Min S, et al. Correction: Compliance with the enhanced recovery after surgery protocol and prognosis after colorectal cancer surgery: A prospective cohort study[J]. *Oncotarget*, 2017, 8(52): 90605
- [19] Svraha M, Wilhelmsen M, Bulut O. Internal hernia following laparoscopic colorectal surgery: single center experience[J]. *Pol Przegl Chir*, 2017, 89(5): 19-22
- [20] Galizia G, Lieto E, Auricchio A, et al. Naples Prognostic Score, Based on Nutritional and Inflammatory Status, is an Independent Predictor of Long-term Outcome in Patients Undergoing Surgery for Colorectal Cancer[J]. *Dis Colon Rectum*, 2017, 60(12): 1273-1284
- [21] Huang J, Xu M, Fang YJ, et al. Association between phytosterol intake and colorectal cancer risk: a case-control study [J]. *Br J Nutr*, 2017, 117(6): 839-850

- [18] 张贤亮,刘勇,刘恩雄,等.复方骨肽注射液辅助治疗肱骨髁间骨折的疗效观察[J].西南国防医药, 2017, 27(8): 868-870
Zhang Xian-liang, Liu Yong, Liu En-xiong, et al. To observe the curative effect of Compound Ossotide Injection adjuvant treatment of intercondylar fracture of humerus [J]. Medical Journal of National Defending Forces in Southwest China, 2017, 27(8): 868-870
- [19] Tsang HHL, Chung HY. The Discriminative Values of the Bath Ankylosing Spondylitis Disease Activity Index, Ankylosing Spondylitis Disease Activity Score, C-Reactive Protein, and Erythrocyte Sedimentation Rate in Spondyloarthritis-Related Axial Arthritis[J]. J Clin Rheumatol, 2017, 23(5): 267-272
- [20] 周贤杰,罗从凤,曾智敏,等.术前 C-反应蛋白和红细胞沉降率对多发骨折术后感染的预测价值[J].中华创伤杂志, 2010, 26(1): 57-60
Zhou Xian-jie, Luo Cong-feng, Zeng Zhi-min, et al. Role of preoperative C-reactive protein and erythrocyte sedimentation rate in predicting postoperative infections following multiple fractures[J]. Chinese Journal of Trauma, 2010, 26(1): 57-60
- [21] 孙晓娜,王晓芳.红细胞沉降率和 C-反应蛋白检测与多发骨折患者术后感染的相关性研究[J].中国伤残医学, 2017, 25(3): 51-53
Sun Xiao-na, Wang Xiao-fang. The study of relationship between infection detection rate and C-reactive protein with multiple fractures in patients with postoperative erythrocyte sedimentation [J]. Chinese Journal of Trauma and Disability Medicine, 2017, 25(3): 51-53
- [22] 刘伟山. 锁定加压钢板内固定术联合骨肽注射液在治疗四肢骨折患者中的疗效观察研究[J].中医临床研究, 2017, 9(11): 114-115
Liu Wei-shan. Observation and study of curative effect of locking plate fixation combined with ossotide injection in the treatment of limbs fracture patients [J]. Clinical Journal of Chinese Medicine, 2017, 9(11): 114-115
- [23] 王斌,邓高鹏,侯平,等.骨肽与复方骨肽治疗中老年四肢骨折的临床效果观察[J].中国老年学杂志, 2014, 34(13): 3764-3765
Wang Bin, Deng Gao-peng, Hou Ping, et al. Clinical effect observation of the fractures in elderly Ossotide and compound Ossotide treatment[J]. Chinese Journal of Gerontology, 2014, 34(13): 3764-3765
- [24] 陈安刚,曾本强,杨勇,等.骨肽注射液促进四肢骨折愈合的效果及对红细胞相关指标影响的研究[J].临床和实验医学杂志, 2013, 12(24): 1963-1965
Chen An-gang, Zeng Ben-qiang, Yang Yong, et al. Ossotide injection to promote the study of limbs fracture healing effect and related indexes of red blood cells [J]. Journal of Clinical and Experimental Medicine, 2013, 12(24): 1963-1965
- [25] 范亮全,徐天波,刘德国,等.骨肽注射液联合接骨七厘片治疗四肢骨折的效果评价[J].中国生化药物杂志, 2017, 37(8): 73-74
Fan Liang-quan, Xu Tian-bo, Liu De-guo, et al. Observation on the effect of bone peptide injection combined with bone grafting in the treatment of limb fractures [J]. Chinese Journal of Biochemical Pharmacology, 2017, 37(8): 73-74
-
- (上接第 1761 页)
- [22] 王公平,杨言通,周博,等.快速康复外科理念应用于胃癌患者围手术期的前瞻性随机对照研究 [J]. 中华胃肠外科杂志, 2014, 17(5): 489-491
Wang Gong-ping, Yang Yan-tong, Zhou Bo, et al. Promotion of postoperative recovery with fast track surgery for gastric cancer patients; undergoing gastrectomy: a prospective randomized controlled study [J]. Chinese Journal of Gastrointestinal Surgery, 2014, 17(5): 489-491
- [23] 韦炳邓,罗中,关秀文,等.快速康复外科对结直肠癌手术效果及炎症因子水平的影响 [J]. 中国现代普通外科进展, 2016, 19(8): 645-648
Wei Bing-deng, Luo Zhong, Guan Xiu-wen, et al. Effect of rapid rehabilitation surgery on the effect of colorectal cancer surgery and the level of inflammatory factors [J]. Chinese Journal of Current Advances in General Surgery, 2016, 19(8): 645-648
- [24] Matsudo T, Ishizaki T, Shigoka M, et al. Examination to Determine Predictive Factors of Effectiveness of Surgical Treatment for Both Pulmonary and Hepatic Metastasis from Colorectal Cancer[J]. Gan To Kagaku Ryoho, 2016, 43(12): 1479-1481
- [25] Back SJ, Kim SH, Kim SY, et al. The safety of a "fast-track" program after laparoscopic colorectal surgery is comparable in older patients as in younger patients[J]. Surg Endosc, 2013, 27(3): 1225-1232
- [26] Coşkun Ö, Öztöpus Ö, Özkan ÖF. Determination of IL-6, TNF- α and VEGF levels in the serums of patients with colorectal cancer [J]. Cell Mol Biol (Noisy-le-grand), 2017, 63(5): 97-101
- [27] Zeng J, Tang ZH, Liu S, et al. Clinicopathological significance of overexpression of interleukin-6 in colorectal cancer [J]. World J Gastroenterol, 2017, 23(10): 1780-1786
- [28] van der Stok EP, Grünhagen DJ, Rothbarth J, et al. The Prognostic Value of Post-operative Serum C-reactive Protein Level for -Survival after Surgery for Colorectal Liver Metastases [J]. Acta Chir Belg, 2015, 115(5): 348-355
- [29] Yang Y, Tang T, Peng W, et al. The comparison of miR-155 with computed tomography and computed tomography plus serum amyloid A protein in staging rectal cancer [J]. J Surg Res, 2015, 193(2): 764-771
- [30] Qi Y, Yao X, Zhang B, et al. Comparison of recovery effect for sufentanil and remifentanil anesthesia with TCI in laparoscopic radical resection during colorectal cancer [J]. Oncol Lett, 2016, 11 (5): 3361-3365