

doi: 10.13241/j.cnki.pmb.2020.18.026

盆底肌功能训练联合阴茎夹对前列腺增生术后患者尿失禁的 临床应用分析*

李雪艳¹ 章宗武² 李令勋^{1△} 李昂灿¹ 许芳¹ 关艳声¹

(1 安徽省第二人民医院泌尿外科 安徽 合肥 230041; 2 安徽医学高等专科学校外科教研室 安徽 合肥 230601)

摘要 目的:分析盆底肌功能训练联合阴茎夹对前列腺增生术后患者尿失禁的临床应用效果。**方法:**选取我院 2017 年 4 月~2019 年 4 月收治的 72 例前列腺增生术后尿失禁患者,随机分为对照组和观察组各 36 例,两组均予盆底肌功能训练,观察组加用阴茎夹控制排尿。对比两组术后尿失禁改善情况、排尿改善情况、国际尿失禁咨询委员会尿失禁问卷表简表(ICI-Q-SF)评分变化、压力性尿失禁分度评价及经济费用情况。**结果:**两组干预后 20 d、干预后 30 d、干预后 90 d 尿失禁发生率均较干预前 10 d 下降,观察组干预后 10 d、干预后 20 d、干预后 30 d、干预后 90 d 尿失禁发生率均低于对照组,差异均有统计学意义($P<0.05$)。两组干预后 90 d 每日总尿量较干预前升高,每日总排尿次数、每日总漏尿次数均较干预前下降;观察组干预后 90 d 每日总尿量高于对照组,每日总排尿次数、每日总漏尿次数均低于对照组,差异均有统计学意义($P<0.05$)。两组干预后 90d ICI-Q-SF 评分均较干预前下降,且观察组干预后 90d ICI-Q-SF 评分低于对照组,差异均有统计学意义($P<0.05$)。观察组患者干预后压力性尿失禁临床治愈率高于对照组,差异有统计学意义($P<0.05$)。两组患者压力性尿失禁分度情况比较差异无统计学意义($P>0.05$)。观察组阴茎夹使用费用为(70.26±8.51)元,低于对照组的(388.71±26.44)元,差异有统计学意义($P<0.05$)。**结论:**在盆底肌功能训练的基础上联合阴茎夹能够有效改善前列腺增生术后患者尿失禁症状及生活质量,且有助于降低患者经济负担,值得临床应用。

关键词:盆底肌功能训练;阴茎夹;前列腺增生;尿失禁;生活质量

中图分类号:R697.3;R694.54 文献标识码:A 文章编号:1673-6273(2020)18-3515-04

Clinical Application of Pelvic Floor Muscle Function Training Combined with Penile Clamp in the Treatment of Urinary Incontinence after Prostatectomy*

LI Xue-yan¹, ZHANG Zong-wu², LI Ling-xun^{1△}, LI Chan-can¹, XU Fang¹, GUAN Yan-sheng¹

(1 Department of Urology Surgery, Second People's Hospital of Anhui Province, Hefei, Anhui, 230041, China;

2 Teaching and Research Office of Surgery, Anhui Medical College, Hefei, Anhui, 230601, China)

ABSTRACT Objective: To analyze the clinical effect of pelvic floor muscle function training combined with penile clamp in the treatment of urinary incontinence after prostatectomy. **Methods:** From April 2017 to April 2019, 72 patients with urinary incontinence after prostatectomy in our hospital were randomly divided into control group and observation group with 36 cases in each group. Both groups were given pelvic floor muscle function training, and the observation group was added with penile clamp to control urination. The improvement of urinary incontinence, the improvement of micturition, the change of ICI-Q-SF score, the grading evaluation of stress urinary incontinence and the economic cost were compared between the two groups. **Results:** The incidence of urinary incontinence in the observation group was lower than that in the control group ($P<0.05$). 90 d after intervention, the total daily urine volume in the two groups was higher than that before intervention, and the total times of urinations per day and total times of leakage of urine per day were lower than that before intervention; the total daily urine volume in the observation group was higher than that in the control group, and the the total times of urinations per day and total times of leakage of urine per day were lower than that in the control group 90d after intervention, with statistical significance ($P<0.05$). The ICI-Q-SF score 90 d after intervention in the two groups was lower than that before intervention, and the ICI-Q-SF score 90d after intervention in the observation group was lower than that in the control group, the difference was statistically significant ($P<0.05$). The clinical cure rate of stress urinary incontinence in the observation group was higher than that in the control group ($P<0.05$), and there was no significant difference in the stress urinary incontinence grade between the two groups ($P>0.05$). The cost of penile clamp in the observation group was (70.26±8.51) yuan, lower than that (388.71±26.44) yuan in the control group, the difference was statistically significant ($P<0.05$). **Conclusion:** On the basis of pelvic floor muscle function training, combined with penile clamp can effectively improve the symptoms of urinary incontinence and quality of life of patients after

* 基金项目:安徽省高等学校自然科学研究项目(12925sk2018b09)

作者简介:李雪艳(1978-),女,硕士研究生,研究方向:泌尿外科,E-mail:13856057422@139.com

△ 通讯作者:李令勋(1969-),博士,主任医师,研究方向:泌尿外科,E-mail:1135087511@qq.com

(收稿日期:2020-03-04 接受日期:2020-03-29)

prostatectomy, and help to reduce the economic burden of patients, which is worthy of clinical application.

Key words: Pelvic floor muscle function training; Penile clamp; Prostatic hyperplasia; Urinary incontinence; Quality of life

Chinese Library Classification(CLC): R697.3; R694.54 **Document code:** A

Article ID: 1673-6273(2020)18-3515-04

前言

良性前列腺增生是中老年男性常见病、高发病;流行病学调查显示,50岁男性良性前列腺增生发病率达40%,80岁以上的老年男性前列腺增生发病率达90%^[1-3]。约有25%的前列腺增生患者因尿路梗阻或合并尿潴留、膀胱结石、肾积水等并发症,需接受手术治疗,但患者术后尿失禁发生率普遍较高^[4-6]。尿失禁的发生不仅会影响患者正常生理功能,还可引发自卑、失落、消极等心理状态变化,降低患者生活质量^[7-8]。因此,控制并改善尿失禁症状也是保证手术效果的关键环节。既往临床常以盆底肌功能训练,效果尚可^[9,10]。近年来,阴茎夹在改善原位膀胱术后尿失禁患者中的积极作用也已得到研究证实^[11]。基于既往研究结论,此次研究就盆底肌功能训练与阴茎夹联合应用于前列腺增生术后尿失禁的效果进行了分析,望为临床尿失禁患者的干预提供新的思路,现报道如下。

1 资料与方法

1.1 一般资料

将2017年4月至2019年4月我院收治的72例前列腺增生术后尿失禁患者纳入此次前瞻性研究。选取标准:(1)明确良性前列腺增生诊断,符合外科手术指征,接受前列腺电切术治疗;(2)术后发生尿失禁,即在拔除导尿管后的不同时间里在未能到达厕所或不能自我控制排尿而有不同程度的尿液漏出;(3)患者对此次研究知情同意,能够配合盆底肌功能训练及阴茎夹使用。排除标准:(1)合并神经源性膀胱、下尿道外伤史伴膀胱出口梗阻;(2)合并严重心肺病变;(3)术前已有尿失禁。使用随机数字法将患者随机分为对照组和观察组各36例,对照组年龄55~79岁,平均(61.28±5.44)岁,病程1周~1个月,平均(11.24±2.35)d;观察组年龄51~83岁,平均(61.59±5.26)岁,病程1周~1个月,平均(12.06±2.44)d。两组患者年龄和病程比较不存在统计学差异($P>0.05$),具有可比性,且本临床研究获得了我院医学伦理委员会授权批准开展实施。

1.2 干预方案

盆底肌功能训练:两组均行盆底肌功能训练。根据患者自身意愿,选择站立位、坐位或平卧位,放松下肢、腹部、臀部肌肉,收紧并上提耻骨、骨尾周围肌肉,行盆底肌自主收缩、肌肉控制训练,维持10s后放松10s,为1组。单次训练循环30组,每日训练3次,30d为1周期,持续训练3个周期。若患者无法掌握训练方法或技巧,则嘱其取侧卧位,戴医用一次性手套并取石蜡油涂于手指,嘱患者放松,缓慢将手指插入肛门,逐步指导患者按上述动作训练,直至患者自觉肛门有明显紧缩感。阴茎夹:对照组常规使用纸尿裤应对尿失禁,观察组给予阴茎夹,在住院期间由患者家属到医疗器械公司自行购买阴茎夹,由医护人员共同指导患者以阴茎夹夹住阴茎根部,定时开放阴茎夹(一般建议每2h开放1次),每次打开阴茎夹排尿后休息

5~10 min,而后重新夹上阴茎夹,以避免阴茎水肿等并发症。

1.3 观察指标

对比两组术后尿失禁改善情况、排尿改善情况、国际尿失禁咨询委员会尿失禁问卷表简表(ICI-Q-SF)评分变化、压力性尿失禁分度评价及经济费用情况。①尿失禁改善情况即干预后10d、干预后20d、干预后30d、干预后90d尿失禁发生率(进行电话随访);②排尿改善情况即干预前、干预后90d每日总排尿量、总排尿次数、总漏尿次数。③ICI-Q-SF评分于干预前、干预后90d进行评估,该量表包括漏尿次数、漏尿量、漏尿对日常生活的影响共3个维度,分值0~21分,得分越高则尿失禁症状越严重^[12]。④压力性尿失禁分度评价^[13]:采用压力性尿失禁标准于干预后90d进行评估。共分为3个度,轻度(仅在咳嗽、喷嚏、大笑等腹压增加情况下发生尿失禁);中度(在屏气、用力、日常活动时或直立、起身站立时发生尿失禁);重度(平躺时发生尿失禁);在便秘、咳嗽等腹压增加情况下也无尿液渗出视为临床治愈。⑤记录两组患者干预后90d内尿失禁物理控制方案的经济费用,即阴茎夹使用费用和纸尿裤使用费用。

1.4 统计学分析

采用SPSS18.0软件进行数据分析,数据的计数资料用(n/%)表示,分析方法采用卡方检验,数据符合正态分布的采用($\bar{x}\pm s$)描述。针对方差齐性的样本,用独立样本T检验分析,否则采用校正T检验处理,等级资料采用秩和检验,以 $P<0.05$ 为差异有统计学意义。

2 结果

2.1 两组患者干预前后尿失禁改善情况

两组干预后20d、干预后30d、干预后90d尿失禁发生率均较干预后10d下降,观察组干预后10d、干预后20d、干预后30d、干预后90d尿失禁发生率均低于对照组,差异均有统计学意义($P<0.05$)。见表1。

2.2 两组患者排尿改善情况比较

两组干预后90d每日总尿量较干预前升高,每日总排尿次数、每日总漏尿次数均较干预前下降;观察组干预后90d每日总尿量高于对照组,每日总排尿次数、每日总漏尿次数均低于对照组,差异均有统计学意义($P<0.05$)。见表2。

2.3 两组患者干预前后ICI-Q-SF评分比较

两组干预后90d ICI-Q-SF评分均较干预前下降,且观察组干预后90d ICI-Q-SF评分低于对照组,差异均有统计学意义($P<0.05$)。见表3。

2.4 两组患者干预后压力性尿失禁分度情况比较

观察组患者干预后压力性尿失禁临床治愈率高于对照组,差异有统计学意义($P<0.05$)。两组患者压力性尿失禁分度情况比较差异无统计学意义($P>0.05$)。见表4。

2.5 两组患者经济费用比较

观察组阴茎夹使用费用为(70.26±8.51)元,低于对照组的

(388.71±26.44)元,差异有统计学意义($T=-68.790, P=0.000$)。

表 1 两组患者干预前后尿失禁发生率比较[n(%)]

Table 1 Comparison of incidence of urinary incontinence between the two groups before and after intervention [n (%)]

Groups	n	10d after intervention	20d after intervention	30d after intervention	90d after intervention
Control group	36	30(83.33)	21(58.33)*	14(38.89)*	10(27.78)*
Observation group	36	19(52.78)	11(30.56)*	6(16.67)*	3(8.33)*
χ^2 value	-	7.730	5.625	4.431	4.600
<i>P</i> value	-	0.005	0.018	0.035	0.032

Note: Compared with 10d after intervention, * $P<0.05$.

表 2 两组患者排尿改善情况比较($\bar{x}\pm s$)Table 2 Comparison of improvement of micturition between the two groups($\bar{x}\pm s$)

Groups	Time	Total daily urine volume	Total times of urinations per	Total times of leakage of
		(mL)	day(time)	urine per day(time)
Control group(n=36)	Before intervention	1025.77±158.42	16.59±2.38	18.41±2.26
	90d after intervention	1339.46±210.59*	11.42±2.01*	13.15±2.10*
Observation group(n=36)	Before intervention	998.64±160.89	16.84±2.60	18.23±1.99
	90d after intervention	1915.40±257.83**	8.51±1.75**	6.41±0.59**

Note: Compared with before intervention, * $P<0.05$; Compared with Control group, ** $P<0.05$.

表 3 两组患者干预前后 ICI-Q-SF 评分比较(分, $\bar{x}\pm s$)Table 3 Comparison of ICI-Q-SF scores between the two groups before and after intervention(score, $\bar{x}\pm s$)

Groups	n	Before intervention	90d after intervention	<i>T</i> value	<i>P</i> value
Control group	36	13.91±2.59	10.26±2.24	6.155	0.000
Observation group	36	14.04±2.33	8.15±1.93	-11.681	0.000
<i>T</i> value	-	-0.223	-4.069	-	-
<i>P</i> value	-	0.412	0.000	-	-

表 4 两组患者干预后尿失禁分度情况比较[n(%)]

Table 4 Comparison of stress urinary incontinence grade between the two groups [n (%)]

Groups	n	Clinical cure	Stress urinary incontinence grade		
			Mild	Moderate	Severe
Control group	36	33(91.67)	3(8.33)	0(0.00)	0(0.00)
Observation group	36	26(72.22)	8(22.22)	1(2.78)	1(2.78)
χ^2/U value	-	4.741		0.043	
<i>P</i> value	-	0.029		2.379	

3 讨论

目前临床对患者术后尿失禁发生的机制尚无明确解释,多数研究认为,术后膀胱功能障碍、膀胱颈纤维化、尿道狭窄及外括约肌损伤,在尿失禁的发生发展中扮演了重要角色^[14-16]。尿失禁的发生不仅会影响手术效果,还可导致患者心理负担增加,甚至引发湿疹、失禁性皮炎等并发症,威胁患者生活质量。因此,及时缓解患者尿失禁症状,对于改善患者预后质量至关重要^[17,18]。男性正常排尿过程需借助膀胱的稳定性和顺应性,膀胱的这两大特性的保障需要借助膀胱颈和前列腺尿道附近的四部分肌肉,即侧括约肌、尿道平滑肌、尿道旁纹肌、尿道外括约

肌,然而当患者进行了前列腺电切术后对膀胱或者临近的侧括约肌造成了一定的损伤,因此患者对排尿过程的控制会造成不同程度的影响,患者排尿过程仅仅只通过外括约肌来进行控制,那么患者在大概率上会出现尿失禁的现象^[19,20]。往常的临床研究中,针对前列腺增生术后尿失禁的病情常采用常采取盆底肌功能训练的方法来进行,通过对盆底肌的锻炼,患者可以在一定程度上提高盆底肌肉的收缩性和张力,从而提高对尿道和膀胱的支撑作用,对促进患者尿道括约肌在短期内恢复张力具有一定的效果,可预防患者术后出现尿失禁的现象^[21-23]。同时,随着尿道括约肌收缩力的改善,患者膀胱顺应性亦可有效恢复,从而延长排尿间隔时长^[24-26]。

本研究对照组接受盆底肌功能训练,其干预后 90 d 尿失禁发生率降至 27.78%, 显现出该方案在控制尿失禁症状的积极作用。同时,得益于盆底肌肌力的恢复,患者尿失禁症状得到有效控制,并表现为排尿的改善及 ICI-Q-SF 评分的显著下降。然而,虽然盆底肌功能训练能够改善远端尿道括约肌循环并加强其功能,减轻创面局部炎症水肿对括约肌关闭机制的影响,但在减少漏尿方面仍存在局限性,患者往往需使用纸尿裤。纸尿裤的应用常伴随着会阴明显不适感,且尿液不良气味对患者日常生活和社会交往的影响较为明显。为解决纸尿裤的弊端,有学者提出阴茎夹方案,即利用阴茎夹的机械控尿原理,通过部分替代尿道括约肌的作用,达到减轻漏尿症状的目的^[27,28]。同时,阴茎夹的使用具有简单方便、经济安全的优势,无需每日定期更换,有助于降低患者经济负担。本研究观察组漏尿症状物理控制费用低于对照组,印证了上述结论。在尿失禁发生率的对比中,可以发现,观察组干预后 10 d、干预后 20 d、干预后 30 d、干预后 90 d 尿失禁发生率均低于对照组,且对两组患者尿失禁分度情况比较,观察组患者临床治愈率显著高于对照组,提示阴茎夹在控制尿失禁症状方面的作用更为明显。另外,与对照组相比,观察组干预后 90 d 每日总尿量有所上升,表明随着尿失禁症状的改善,患者对漏尿症状的畏惧心理减轻,每日水分摄入量有所增加,该结果也侧面反映了阴茎夹在改善尿失禁患者心理状态方面的积极作用。国外研究表明,阴茎夹在尿失禁的治疗中安全、有效,且能够被多数患者所接受^[29,30],本研究结果显示,观察组干预后 90 d ICI-Q-SF 评分低于对照组,说明其生活质量改善更为明显,在印证上述结论的同时,也为阴茎夹的推广应用奠定了坚实的基础。

综上所述,在盆底肌功能训练的基础上联合阴茎夹能够有效改善前列腺增生术后患者尿失禁症状及生活质量,且有助于降低患者经济负担,值得临床推广应用。

参考文献(References)

- 袁冰,张金龙,王茂强,等.前列腺动脉栓塞术治疗巨大前列腺增生伴严重肾功能损伤一例[J].中华医学杂志,2019,99(25):1991-1992
- 詹长生,曹乃龙,王啸虎,等.前列腺动脉栓塞术治疗前列腺增生的研究进展和现状[J].山西医科大学学报,2018,49(4):434-438
- 朱自强.良性前列腺增生最新研究进展[J].临床泌尿外科杂志,2019,34(5):409-412
- Hirasawa Y, Kato Y, Fujita K. Age and prostate volume are risk factors for transient urinary incontinence after transurethral enucleation with bipolar for benign prostatic hyperplasia [J]. Int J Urol, 2018, 25(1): 76-80
- Tienza A, Hevia M, Merino I, et al. Can low urinary tract symptoms influence postprostatectomy urinary incontinence?[J]. Minerva Urol Nefrol, 2016, 68(4): 324-329
- 杨海,唐贤富,徐小涵,等.保留前列腺尖部尿道黏膜的经尿道前列腺电切术改善良性前列腺增生术后尿失禁的临床研究[J].微创泌尿外科杂志,2019,8(3):194-199
- Petrillo M, Pesapane F, Fumarola EM, et al. State of the art of prostatic arterial embolization for benign prostatic hyperplasia [J]. Gland Surg, 2018, 7(2): 188-199
- Kretschmer A, Gratzke C. Therapeutic options for stress urinary incontinence following surgery for benign prostate enlargement [J]. Aktuelle Urol, 2018, 49(4): 334-338
- 黄月娥,赵格格.针灸联合盆底肌群康复训练对前列腺增生电切患者术后尿失禁与勃起功能障碍发生率的影响 [J]. 中国性科学, 2019, 28(4): 113-116
- 洪媚,宋双临,谢奎,等.针灸配合盆底肌功能训练治疗良性前列腺增生术后压力性尿失禁疗效观察 [J]. 上海针灸杂志, 2020, 39(2): 206-211
- 全翔凤,姚莉,唐冬媛,等.阴茎夹在原位膀胱术后尿失禁男性病人中的应用效果[J].护理研究, 2015, (33): 4156-4157
- 邵魁卿,高瞻,沈建武,等.芪实颗粒联合盆底肌训练对女性轻中度压力性尿失禁患者尿失禁问卷量表评分、尿失禁生活质量量表评分的影响[J].中国医院用药评价与分析, 2019, 19(6): 696-698, 701
- 邓丽群,马苏亚.盆底超声在压力性尿失禁中的临床应用及进展[J].现代实用医学, 2018, 30(11): 1419-1422
- chaake W, van der Schaaf A, van Dijk LV, et al. Development of a prediction model for late urinary incontinence, hematuria, pain and voiding frequency among irradiated prostate cancer patients[J]. PLoS One, 2018, 13(7): e0197757
- Tantawy SA, Elgohary HMI, Abdelbasset WK, et al. Effect of 4 weeks of whole-body vibration training in treating stress urinary incontinence after prostate cancer surgery: a randomised controlled trial[J]. Physiotherapy, 2019, 105(3): 338-345
- Albkri A, Girier D, Mestre A, et al. Urinary Incontinence, Patient Satisfaction, and Decisional Regret after Prostate Cancer Treatment: A French National Study[J]. Urol Int, 2018, 100(1): 50-56
- Fujimura T, Igawa Y, Aizawa N, et al. Longitudinal change of comprehensive lower urinary tract symptoms and various types of urinary incontinence during robot-assisted radical prostatectomy [J]. Neurourol Urodyn, 2019, 38(4): 1067-1075
- Cozzarini C, Rancati T, Palorini F, et al. Patient-reported urinary incontinence after radiotherapy for prostate cancer: Quantifying the dose-effect[J]. Radiother Oncol, 2017, 125(1): 101-106
- 邓丽群,马苏亚.盆底超声在压力性尿失禁中的临床应用及进展[J].现代实用医学, 2018, 30(11): 1419-1422
- 李素琼,刘翔宇,肖川,等.经尿道前列腺激光汽化电切术后发生尿失禁危险因素分析及应对措施[J].实用医院临床杂志, 2019, 16(5): 153-156
- Davenport MH, Nagpal TS, Mottola MF, et al. Prenatal exercise (including but not limited to pelvic floor muscle training) and urinary incontinence during and following pregnancy: a systematic review and meta-analysis[J]. Br J Sports Med, 2018, 52(21): 1397-1404
- Woodley SJ, Boyle R, Cody JD, et al. Pelvic floor muscle training for prevention and treatment of urinary and faecal incontinence in antenatal and postnatal women[J]. Cochrane Database Syst Rev, 2017, 12(12): CD007471
- Casey EK, Temme K. Pelvic floor muscle function and urinary incontinence in the female athlete [J]. Phys Sportsmed, 2017, 45(4): 399-407
- Kahyaoglu Sut H, Balkanli Kaplan P. Effect of pelvic floor muscle exercise on pelvic floor muscle activity and voiding functions during pregnancy and the postpartum period[J]. Neurourol Urodyn, 2016, 35(3): 417-22

- Hematoma Associated with Spontaneous Intracranial Hypotension: Therapeutic Strategies and Outcomes of 55 Cases [J]. *Neurologia Medico Chirurgica*, 2015, 56(2): 69
- [16] M Dobran, M Iacoangeli, A R Scortichini, et al. Spontaneous chronic subdural hematoma in young adult: the role of missing coagulation facto[J]. *IL Giornale Di Chirurgia*, 2017, 38(2): 66-70
- [17] Katsigiannis S, Hamisch C, Krischek B, et al. Independent predictors for functional outcome after drainage of chronic subdural hematoma identified using a logistic regression model [J]. *Journal of neurosurgical sciences*, 2020, 64 (2): 133-140
- [18] Marshman L. Letter to the editor Re: Standardized assessment of outcome and complications in chronic subdural hematoma: results from a large case series by Bucher B [J]. *Acta neurochirurgica*, 2019, 161(12): 2483-2484
- [19] Andersen-Ranberg NC, Debrabant B, Poulsen FR, et al. The Danish chronic subdural hematoma study-predicting recurrence of chronic subdural hematoma[J]. *Acta neurochirurgica* 2019, 161(5): 885-894
- [20] Fiorella D, Arthur AS. Middle meningeal artery embolization for the management of chronic subdural hematoma [J]. *Journal of neurointerventional surgery*, 2019, 11(9): 912-915
- [21] Gun Seok Lee, Young Seok Park, Kyung Soo Min, et al. Spontaneous Resolution of a Large Chronic Subdural Hematoma Which Required Surgical Decompression[J]. *Journal of Korean Neurosurgical Society*, 2015, 58(3): 301
- [22] Motiei-Langroudi R, Thomas AJ, Ascanio L, et al. Factors Predicting the Need for Surgery of the Opposite Side After Unilateral Evacuation of Bilateral Chronic Subdural Hematomas[J]. *Neurosurgery*, 2019, 85 (5): 648-655
- [23] Rosario Maugeri, Antonella Giugno, Francesca Graziano, et al. Delayed chronic intracranial subdural hematoma complicating resection of a tanycytic thoracic ependymoma[J]. *Surgical Neurology International*, 2016, 7(2): 20
- [24] Tange M, Yoshida M, Nakai Y, et al. The Role of an Impurity in Ceftriaxone Sodium Preparation for Injection in Determining Compatibility with Calcium-Containing Solutions [J]. *Chemical & pharmaceutical bulletin*, 2016, 64 (3): 207-14
- [25] Duan HY, Liu DM, Qian P, et al. Effect of atorvastatin on plasma NT-proBNP and inflammatory cytokine expression in patients with heart failure [J]. *Genetics and molecular research : GMR*, 2015, 14 (4): 15739-48
- [26] Yong H, Wang X, Mi L, et al. Effects of atorvastatin loading prior to primary percutaneous coronary intervention on endothelial function and inflammatory factors in patients with ST-segment elevation myocardial infarction [J]. *Experimental and therapeutic medicine*, 2014, 7(2): 316-322
- [27] David Yuen Chung Chan, Danny Tat Ming Chan, Tin Fung David Sun, et al. The use of atorvastatin for chronic subdural haematoma: a retrospective cohort comparison study [J]. *British Journal of Neurosurgery*, 2016, 31(1): 1-6
- [28] Mayyas F, Baydoun D, Ibdah R, et al. Atorvastatin Reduces Plasma Inflammatory and Oxidant Biomarkers in Patients With Risk of Atherosclerotic Cardiovascular Disease [J]. *Journal of cardiovascular pharmacology and therapeutics*, 2018, 23(3): 216-225
- [29] Lang Y, Hongying M I, Liu M, et al. Interventional effects of tongxinluo combined with atorvastatin and aspirin on adventitial inflammation in early stage of atherosclerosis [J]. *Chinese Journal of Pathophysiology*, 2015, 31(1): 154-159
- [30] Walker M E, Souza P R, Colas R A, et al. 13-series resolvins mediate the leukocyte-platelet actions of atorvastatin and pravastatin in inflammatory arthritis [J]. *Faseb Journal Official Publication of the Federation of American Societies for Experimental Biology*, 2017, 31 (8): 3636
- [31] Maria Licci, Maria Kamenova, Raphael Guzman, et al. Influence of Postoperative Thrombosis Prophylaxis on the Recurrence of Chronic Subdural Hematoma After Burr-Hole Drainage [J]. *Critical Care Medicine*, 2017, 46(1): 1
- [32] Xunhui Yuan, Xiaodong Shi, Hang Xiao, et al. Intracranial Subdural Empyema Mimicking Chronic Subdural Hematoma [J]. *Journal of Craniofacial Surgery*, 2016, 27(2): 1
- [33] Ratender Kumar Singh, Vikas Agarwal, Arvind Kumar Baronia, et al. The Effects of Atorvastatin on Inflammatory Responses and Mortality in Septic Shock: A Single center, Randomized Controlled Trial [J]. *Indian J Crit Care Med*, 2017, 21(21): 646-54
- [34] Perez-Calahorra S, Laclaustra M, Marco-Benedi V, et al. Comparative efficacy between atorvastatin and rosuvastatin in the prevention of cardiovascular disease recurrence [J]. *Lipids in health and disease*, 2019, 18(1): 216
- [35] Tian J, Tai Y, Shi M, et al. Atorvastatin Relieves Cognitive Disorder After Sepsis Through Reverting Inflammatory Cytokines, Oxidative Stress, and Neuronal Apoptosis in Hippocampus [J]. *Cellular and molecular neurobiology*, 2020, 40(4): 521-530

(上接第 3518 页)

- [25] Aydın Sayılan A, Özbaş A. The Effect of Pelvic Floor Muscle Training On Incontinence Problems After Radical Prostatectomy[J]. *Am J Mens Health*, 2018, 12(4): 1007-1015
- [26] Oliveira M, Ferreira M, Azevedo MJ, et al. Pelvic floor muscle training protocol for stress urinary incontinence in women: A systematic review[J]. *Rev Assoc Med Bras (1992)*, 2017, 63(7): 642-650
- [27] Koifman L, Hampl D, Silva MI, et al. Treatment Options and Outcomes of Penile Constriction Devices[J]. *Int Braz J Urol*, 2019, 45 (2): 384-391
- [28] Levy A, Fader M, Bader D, et al. Penile compression clamps: A model of the internal mechanical state of penile soft tissues [J]. *Neurourol Urodyn*, 2017, 36(6): 1645-1650
- [29] Sharaf A, Fader M, Macaulay M, et al. Use of an occlusive penile clamp during filling cystometry in men with symptoms of stress urinary incontinence [J]. *Low Urin Tract Symptoms*, 2019, 11 (3): 133-138
- [30] Nazarko L. Male urinary incontinence management: penile sheaths [J]. *Br J Community Nurs*, 2018, 23(3): 110-116