

人外周血树突状细胞体外诱导扩增的研究

吴楠¹ 刘薇² 卢光琇^{2△}

(1 深圳南山区疾病预防控制中心 广东 深圳 518054 ;2 中南大学干细胞研究所 湖南 长沙 410078)

摘要 目的 探讨可用于临床治疗功能成熟的 DC 体外扩增的优化培养方案。方法 胎牛血清培养基联合细胞因子 rhGM-CSF(100 ng/mL)和 rhIL-4(50 ng/mL)扩增人外周血分离的单个核细胞,细胞培养分别按 $5 \times 10^6/\text{mL}$ 、 $6 \times 10^6/\text{mL}$ 和 $7 \times 10^6/\text{mL}$ 的密度,加入 6 孔培养板。第 6 d 加入 rhTNF- α (100 ng/mL)联合培养,分别于第 6 d,第 9 d 和 12 d 收获细胞。从形态学、细胞表面标志方面进行鉴定。结果 显微镜观察,经过 9 d 诱导后,培养细胞具有典型树突状细胞外形。流式细胞仪分析, $6 \times 10^6/\text{mL}$ 密度的细胞培养组培养到第 9 天最宜。结论 细胞具有典型的 DC 的形态特征,细胞表型及功能实验证实其 DC 的特性,说明建立的血清培养基联合细胞因子 rhGM-CSF、rhIL-4 和 rhTNF- α 体外诱导 DC 的方法是切实可行的。

关键词 树突状细胞 细胞培养 细胞因子

中图分类号 R392.12, Q813, Q42 **文献标识码** A **文章编号** 1673-6273 (2012) 25-4948-02

Experimental Study on Induction and Expansion of Dendritic Cells from Human Peripheral Blood In Vitro

WU Nan¹, LIU Wei², LU Guang-xiu^{2△}

(1 Shenzhen NanShan Centre for Disease Control and Prevention, Guangdong, Shenzhen 518054, China;

2 Laboratory of stem cell, Center south university, Changsha, 410078, China)

ABSTRACT Objective: To investigate a culture system for DC expansion. **Methods:** Using serum medium with rhGM-CSF (100ng/mL) and rhIL-4 (50ng/mL) for 6 days, rhTNF- α (100ng/mL) was added on the 6th day of culture. Human peripheral blood were seeding at the concentration of $5 \times 10^6/\text{mL}$, $6 \times 10^6/\text{mL}$ and $7 \times 10^6/\text{mL}$ in 6-well flat-bottomed plates. Suspension cells were harvested at days 6, 9 and 12 and analysed on morphology, phenotype observed. **Result:** Cells on day 9 of culture displayed DC-like morphology. Phenotype analysis on DCs acquired showed that the cells of $6 \times 10^6/\text{mL}$ cultured for 9 days contained a higher percentage of DC cells. **Conclusion:** The results suggested the culture method was feasible.

Key words: Dendritic cell; Cell Culture; Cytokine

Chinese Library Classification(CLC): R392.12, Q813, Q42 **Document code:** A

Article ID: 1673-6273 (2012) 25-4948-02

前言

树突状细胞(Dendritic cells, DCs)是 1973 年 Steinman 和 Cohn 在小鼠脾脏贴壁细胞中发现的一种新型细胞,因其表面具有星状多形性或树枝状突起而得名^[1]。DC 因其表面具有大量的 MHC 分子和共刺激分子而具有递呈抗原的能力,激发机体的免疫应答,也因此而成为肿瘤免疫治疗的手段之一。然而 DC 数量极少,约占外周血单个核细胞的 1%,且广泛分布于全身淋巴组织和非淋巴组织中,直接从组织中分离、提取十分困难。DC 体外培养及扩增是 DC 应用于临床的前提条件。本研究旨在探索 DC 体外扩增的优化培养方案。

1 材料和方法

1.1 实验材料

细胞因子 rhGM-CSF 购自美国 R&D Systems 公司,细胞因子 rhIL-4 购自美国 R&D Systems 公司,细胞因子 rhTNF- α 购自

美国 R&D Systems 公司,胎牛血清培养基购自中国四季青公司。

1.2 方法

1.2.1 人外周血树突状细胞的培养 采用密度梯度离心法分离外周血单个核细胞。将洗涤后的细胞重悬于胎牛血清培养液中,并加入细胞因子 rhGM-CSF 100 ng/mL, rhIL-4 50 ng/mL 调细胞密度为 $5 \times 10^6/\text{mL}$ 、 $6 \times 10^6/\text{mL}$ 和 $7 \times 10^6/\text{mL}$ 加入 6 孔板每孔 2 mL,置于 37℃,5% CO₂ 培养箱。培养第 6 d,换液时同时加入细胞因子 rhTNF- α 100 ng/mL,继续培养。

1.2.2 人外周血树突状细胞的鉴定 分别于第 6 d、9 d、12 d 收集悬浮细胞,采用流式细胞仪对细胞表面分子 CD1a 鉴定。

1.2.3 统计分析方法 以 Epidata3.0 建立数据库,采用 SPSS 13.0 进行分析,所有数据采用均数 + 标准差表示。

2 结果

2.1 树突状细胞形态观察

从外周血获得单个核细胞,贴壁 2h 后,去悬浮细胞,获得的贴壁细胞即单核细胞。加入细胞因子 rhGM-CSF 100 ng/mL, rhIL-4 50 ng/mL。培养到第 6 d 加入细胞因子 rhTNF- α 100 ng/mL,培养到第 9 d 可见大部分细胞悬浮不贴壁,胞质变大,

作者简介 吴楠(1980),男,博士,研究方向 肿瘤免疫。

电话 0755-23641257, E-mail: wnwunan@163.com

△通讯作者 卢光琇, E-mail: cslgx@126.com

(收稿日期 2012-05-11 接受日期 2012-06-08)

形态不规则。培养板上仅有少量贴壁,形态拉长的巨噬细胞(图1)。




图1 第9天细胞形态

Fig.1 The morphology of cell at 9 day

2.2 细胞培养密度和培养时间的细胞表型比较

细胞培养密度 $5 \times 10^6/\text{mL}$, $6 \times 10^6/\text{mL}$ 和 $7 \times 10^6/\text{mL}$ 分别于第6 d, 9 d, 12 d 通过流式细胞仪检测 DC 特异的细胞表面标志分子 CD1a, 发现细胞密度 $6 \times 10^6/\text{mL}$ 和培养到9 d 时, 培养 DC 的效果最好(表1)。

表1 细胞表面标志 CD1a 分析

Table 1 Analysis of Phenotype CD1a of cells

Culture density	Culture time		
	At 6 day CD1a(%)	At 9 day CD1a(%)	At 12 day CD1a(%)
$5 \times 10^6/\text{mL}$	10.83 + 1.50	30.69 + 4.42	16.25 + 2.81
$6 \times 10^6/\text{mL}$	37.15 + 2.72	65.00 + 5.27	27.05 + 0.52
$7 \times 10^6/\text{mL}$	34.43 + 0.53	46.95 + 1.36	19.85 + 1.74

综合培养密度和培养时间考虑, 以细胞密度 $6 \times 10^6/\text{mL}$ 和培养时间9天为宜。本研究的 DC 培养方案为进一步研究及临床应用打下了基础。

参考文献 (References)

[1] Steinman R, Cohn Z. Identification of a novel cell type in peripheral lymphoid organs of mice[J]. J Exp Med, 1973,137(3):1142

[2] Gavanagh L,Von-Andrian H. Travellers in many guises: the origins and destinations of dendritic cells [J]. Immunol Cell Bio, 2002,80(5): 448-462

[3] Zehn D, Cohn C J. Extended presentation of specific MHC-peptide complexes by mature dendritic cells compared to other types of antigen-presenting cells[J]. Eur J Immunol, 2004,34(6):1551-1560

[4] Chung NP, Chen Y, Chan VS, et al. Dendritic cells: sentinels against pathogens[J]. Histol his to pathol, 2004,19 (1): 317-324

[5] Brossart P, Wirths S, Brugger W, et al. Dendritic cells in cancer vaccines[J]. Exp Hernatol, 2001,29(11):1247-1255

[6] Morel PA, Feili-Hariri M,Coates PT, et al. Dendritic cells, T cell tolerance and therapy of adverse immunere actions [J]. Clin Exp Immunol,2003,133(1):1-10

[7] Lijun Z, Xin Z, Danhua S, et al. Tumor-infiltrating dendritic cells may

3 讨论

DC 起源于骨髓多功能造血干细胞^[2],是迄今发现的功能最强的抗原呈递细胞(antigen-presenting cell,APC),其激发 T 细胞增殖能力及抗原呈递能力是巨噬细胞和 B 细胞的 100~1000 倍^[3]。而且 DC 是体内唯一能够活化 Naive T 细胞的 APC,并能特异性的激活细胞毒性 T 细胞,在抗感染、抗肿瘤和治疗自身免疫性疾病等过程中发挥了重要作用^[4-8]。然而,尽管 DC 在体内广泛分布于除脑以外的各个脏器^[9-11],但数量少(仅占外周血单个核细胞的 1%左右)且分散,不能满足体内或体外研究及临床应用的需要。因而在体外利用前体细胞对未成熟 DC 进行扩增有重大意义^[12]。

DC 表面表达丰富的免疫分子,不同种属的 DC 其细胞表面特异性标志物并不相同^[13]。CD1a 是目前国内外鉴定人 DC 常用的主要分子。CD1a 主要表达于人胸腺细胞、DC 细胞表面,是鉴定 DC 的最好标记^[14-18]。本研究在培养时发现初始细胞的浓度对于集落的形成也很重要。我们对不同培养细胞密度比较发现培养细胞密度 $6 \times 10^6/\text{mL}$, 使 DC 前体细胞能互相接触有利于集落的形成但又不至于太拥挤互相抑制生长。我们对不同的培养时间比较发现培养到第9天,培养细胞其表面共刺激分子 CD1a 的表达率最高。

be used as clinicopathologic prognostic factors in endometrial carcinoma[J]. Int J Gynecol Cancer, 2012,22(5):836-841

[8] Chen Z, Yu B, Wu XL, et al. Carboxymethylpachymaran enhances immunologic function of dendritic cells cultured in two kinds of hepatoma carcinoma cell line's supernatant via nuclear factor κ B/Rel pathway[J]. Chin J Integr Med, 2012,18(3):203-208

[9] Banchereau J, Briere F, Caux C, et al. Immunobiology of dendritic cells[J]. Annu Rev Immunol, 2000, 18(4) :767-811

[10] Segura E, Valladeau-Guilemond J, et al. Characterization of resident and migratory dendritic cells in human lymph nodes [J]. J Exp Med, 2012,209(4):653-660

[11] Jiangyang L, Qian L, Donnadieu MH, et al. Changes of spleen dendritic cells in the terminal stage of multiple organ dysfunction syndrome[J]. Acta Biomed, 2011,82(2):146-153

[12] Ning J, Morgan D, Pamphilon D, et al. A rapid culture technique produces functional dendritic-like cells from human acute myeloid leukemia cell lines[J]. J Biomed Biotechnol, 2011,5(4)123-132

[13] Shortman K, Liu YJ. Mouse and human dendritic cells subtypes[J]. Nat Rev Immunol, 2002,2(3):151-161

(下转第 4997 页)

要密切监测肝、肾功能,根据病情及时调整剂量,把肝损害的危险程度降至最低;尽量不要与其他药物合用,避免药物相互作用。由于儿童用药不能进行临床试验,所以对儿童用药一定需要特别谨慎。

参考文献 (References)

- [1] Koval'chuk VV, Efimov MA. Efficacy and tolerability of short courses of nonsteroid anti-inflammatory drugs in the treatment of dorsalgia: results of the comparative study [J]. Zh Nevrol Psikhiatr Im S S Korsakova, 2010,110(1): 55-58
- [2] Kean W.F, Buchanan W.W. The use of NSAIDs in rheumatic disorders 2005: a global perspective [J]. Inflammopharmacology, 2005, 13: 343-370
- [3] Chopra S, Saini RK, Sanyal SN. Intestinal toxicity of non-steroidal anti-inflammatory drugs with differential cyclooxygenase inhibition selectivity [J]. Nutr Hosp, 2007, 22(5): 528-537
- [4] Ilic KV, Sefik-Bukilica M, Jankovic S. Efficacy and safety of two generic copies of nimesulide in patients with low back pain or knee osteoarthritis [J]. Reumatismo, 2009, 61(1): 27-33
- [5] Su B, Cai X, Hong Y. COX-2 inhibitor nimesulide analogs are aromatase suppressors in breast cancer cells [J]. J Steroid Biochem Mol Biol, 2010, 122(4):232-238
- [6] Xu JM. Therapeutic effect and adverse reaction observation of Nimesulide to hyperpyrexia child [J]. Public Medical Forum Magazine, 2009, 13: 31-32
- [7] Piyush Gupta, HPS. Sachdev. Safety of oral use of Nimesulide in children: systematic review of randomized controlled trials [J]. Indian Pediatrics, 2003, 40:518-531
- [8] Wang YL. Pharmacologic actions and pharmacologic actions of Nimesulide [J]. China contemporary Medicine, 2006, 3(5):52-53
- [9] Candelario-Jalil E. Nimesulide as a promising neuroprotectant in brain ischemia: new experimental evidences [J]. Pharmacol Res, 2008, 57(4):266-273
- [10] Laporte JR., Ibanez I, Vidal X. Upper gastro-intestinal bleeding associated with the use of NSAIDs: newer vs. older agents [J]. Drug Safety, 2004, 27: 411-420
- [11] Bjarnason I, Thjodleifsson B. Gastrointestinal toxicity of non-steroidal anti-inflammatory drugs: the effect of nimesulide compared with naproxen on the human gastrointestinal tract [J]. Rheumatology, 1999, 38: 24-32
- [12] Zhang L, Fan CH. Nimesulide induced one liver functional lesion and upper gastrointestinal hemorrhage [J]. Chinese Journal of Drug Application and Monitoring, 2007, (2): 63-64
- [13] Venegoni M., Da Cas R, Menniti-Ippolito F. Effects of the European restrictive actions concerning nimesulide prescription: a simulation study on hepatopathies and gastrointestinal bleedings in Italy [J]. Ann Ist Super Sanita, 2010, 46(2):153-157
- [14] Bessone F. Non-steroidal anti-inflammatory drugs: What is the actual risk of liver damage [J]. World J Gastroenterol, 2010, 16 (45): 5651-5661
- [15] Yu RH. Retrospective analysis of Nimesulide cause liver severity adverse reaction [J]. China Medical Herald, 2008, 5(9): 175-175
- [16] Zhu JX, Shi QP, Xing R, et al. Literature analysis on the preparations for oral of nimesulide induced adverse reactions [J]. China Medical Herald, 2008, 5(36): 104-105
- [17] He P, Zhang SR., Qiang DM., et al. Analysis of nimesulide induced five neonatus kidney adverse reaction [J]. Shaanxi Medical Journal, 2009, 38(10): 1410-1410
- [18] Khanna D, Khanna PP, Furst DE. COX-2 controversy: where do we go from here [J]. Inflammopharmacology, 2005, 13: 395-402
- [19] Carrasco-Portugal Mdel C, Flores-Murrieta FJ. Comparison of suspension composition on the pharmacokinetics of nimesulide in rats [J]. Proc West Pharmacol Soc, 2008, 51: 58-59
- [14] Coventry BJ, Austyn JM Chrysidsis S, et al. Identification and isolation of CD1a positive putative tumour infiltrating dendritic cells in human breast cancer [J]. Adv Exp Med Biol, 1997,417(1):571-577
- [15] Anjana R, Joseph L, Suresh R. Immunohistochemical localization of CD1a and S100 in gingival tissues of healthy and chronic periodontitis subjects [J]. Oral Dis, 2012,4(3):160-164
- [16] Oztürk A, Yildiz L. Expression of transient receptor potential vanilloid receptor 1 and toll-like receptor 4 in aggressive periodontitis and in chronic periodontitis [J]. J Periodontal Res, 2011,46(4):475-482
- [17] Winning T, Gemmell E, Polak B, et al. Expression of CD1a monocytes cultured with supernatants from periodontally diseased gingival epithelial cells [J]. Oral Dis, 1996,2(4):247-252
- [18] Athanasas-Platsis S, Savage NW, et al. Induction of the CD1a Langerhans cell marker on human monocytes [J]. Arch Oral Biol, 1995,40(2): 157-160

(上接第 4949 页)