

doi: 10.13241/j.cnki.pmb.2017.10.036

## 七氟醚联合骶管阻滞麻醉对小儿疝气手术的麻醉效果分析

高洁 毕翻利<sup>△</sup> 党璐 何秀莉 胡彬

(延安大学附属医院麻醉科 陕西 延安 716000)

**摘要 目的:**观察七氟醚联合骶管阻滞麻醉对小儿疝气手术的麻醉效果。**方法:**选取 80 例行腹股沟疝气手术患儿,按随机数字表法分为两组,对照组(39 例)静脉注射氯胺酮,观察组(41 例)先吸入 8%七氟醚,然后进行骶管阻滞麻醉,通过观察并记录两组患儿生命体征、麻醉诱导时间、苏醒时间、手术麻醉时间、苏醒期躁动评分(Pediatric anesthesia emergence delirium, PAED)和麻醉诱导期合作量表(Induction Compliance Checklist, ICC)及麻醉期间不良反应情况,评价七氟醚联合骶管阻滞麻醉对小儿疝气手术的麻醉效果。**结果:**两组切皮后 T1、T2 时心率(HR)、平均动脉压(MAP)水平均高于 T0 时的值( $P<0.05$ ),两组切皮后 T1、T2 时组间 HR、MAP 水平相比,无统计学差异( $P>0.05$ )。两组切皮后 T3 时 HR、MAP 水平基本恢复到 T0 时的水平。两组切皮前后 4 个时间点的血氧饱和度( $SpO_2$ )相比,无统计学差异( $P>0.05$ )。观察组患儿麻醉诱导时间,苏醒时间均明显短于对照组患儿( $P<0.05$ ),两组术中麻醉持续时间相比,无统计学差异( $P>0.05$ ),均能达到期望麻醉时间,观察组患儿 PAED 评分和 ICC 评分均低于对照组患儿( $P<0.05$ ),不良反应组间比较无统计学差异( $P>0.05$ )。**结论:**七氟醚联合骶管阻滞麻醉对小儿疝气手术具有良好的麻醉效果,麻醉诱导快,苏醒快,小儿配合度高,术后躁动少,值得临床推广使用。

**关键词:**七氟醚;骶管阻滞麻醉;麻醉诱导时间;苏醒期躁动;不良反应

**中图分类号:**R726; R614 **文献标识码:**A **文章编号:**1673-6273(2017)10-1934-04

## Anesthetic Effect of Sevoflurane Combined with Sacral Block Anesthesia in Treatment of Pediatric Hernia Surgery

GAO Jie, BI Fan-li<sup>△</sup>, DANG Lu, HE Xiu-li, HU Bin

(Department of Anesthesiology, Yanan University Affiliated Hospital, Yan'an, Shaanxi, 716000, China)

**ABSTRACT Objective:** To discuss the anesthetic effect of sevoflurane combined with sacral block anesthesia in treatment of pediatric hernia surgery. **Methods:** 80 children with inguinal hernia surgery were selected, they were divided into two groups randomly. The control group (39 cases) was given ketamine by intravenous injection. The observation group (41 cases) was inhaled 8 % sevoflurane, then given sacral block anesthesia. The anesthetic effect of sevoflurane combined with sacral block anesthesia in treatment of pediatric hernia surgery was evaluated by vital sign, induction time, recovery time, anesthesia time, Pediatric anesthesia emergence delirium (PAED). Induction Compliance Checklist (ICC) score and adverse reactions during anesthesia. **Results:** At T1 and T2, the HR, MAP of two groups were higher than that of the T0 ( $P<0.05$ ). But there were no statistical significance on HR and MAP at T1 and T2 ( $P>0.05$ ). At T3, the level of HR, MAP was close to the value of T1. At 4 point of time, there were no statistical significance on  $SpO_2$  between two groups ( $P>0.05$ ). The induction time and recovery time of observation group was shorter than that of the control group ( $P<0.05$ ). There were no statistical significance on anesthesia duration between two groups ( $P>0.05$ ). The expected anesthesia time was achieved in two groups. The PAED and ICC scores of observation group were lower than that of the control group ( $P<0.05$ ). There were no statistical significance on the incidence of adverse reactions between two groups ( $P>0.05$ ). **Conclusions:** Sevoflurane anesthesia combined with sacral block anesthesia have good anesthesia effect on pediatric hernia surgery with short anesthesia induction, and the children can recover quickly and cooperate well with less postoperative agitation, worthy of clinical use.

**Key words:** Sevoflurane; Sacral block anesthesia; Anesthesia induction time; Restlessness in recovery period; Adverse reaction

**Chinese Library Classification(CLC):** R726; R614 **Document code:** A

**Article ID:** 1673-6273(2017)10-1934-04

### 前言

腹股沟疝是小儿外科常见的疾病,多采用小儿疝囊高位结

扎术进行治疗。氯胺酮静脉麻醉为小儿常用的麻醉方法,但氯胺酮可导致患儿心率增快、苏醒延迟、术后躁动,再加上小儿对针刺性麻醉方式抗拒性强,因此临床需选择更安全有效、患儿顺应性高的麻醉方式<sup>[1]</sup>。七氟醚为吸入性麻醉药物,麻醉效果好,苏醒快。小儿生理特点与成人不同,对麻醉药物耐受性更强,麻醉处理存在一定难度<sup>[2]</sup>。而采用骶管阻滞麻醉可较好地解决这一问题。既往研究中,麻醉药物及麻醉方式对患儿生命体征、麻醉效果的影响多有报道<sup>[3,4]</sup>,然而小儿麻醉尤其需要注意

作者简介:高洁(1983-),女,本科,主治医师,研究方向:临床麻醉,电话:13772881067, E-mail: gaojie\_1983@medicpaper.cn

<sup>△</sup> 通讯作者:毕翻利(1983-),女,本科,主治医师,研究方向:临床麻醉,电话:18091106126, E-mail: bifanli\_1983@medicpaper.cn

(收稿日期:2016-09-21 接受日期:2016-10-20)

安全性问题,为此,本文回顾性分析 2014 年 3 月 -2016 年 3 月选取的 80 例需行小儿疝气手术的患儿,分别用七氟醚联合骶管阻滞麻醉和静脉注射氯胺酮进行麻醉,并对麻醉效果及安全性进行比较,对苏醒期躁动、小儿麻醉诱导时的行为表现也进行了评价。现报道如下。

## 1 资料与方法

### 1.1 病例资料

选取腹股沟疝气手术患儿 80 例,年限:2014 年 3 月 -2016 年 3 月,纳入标准:①术前诊断为单侧腹股沟疝;②美国麻醉医师协会(ASA)分级为 I - II 级;③患儿年龄 2~6 岁;④经本院伦理委员会同意,术前患儿家属均签署书面知情同意书。排除标准:合并肝肾功能异常、心脑血管疾病的患儿。按随机数字表法将患儿分为两组,对照组(39 例)患儿术前静脉注射氯胺酮,其中,男 25 例,女 14 例,平均年龄( $5.9 \pm 1.5$ )岁;观察组(41 例)术前吸入 8%的七氟烷,然后行骶管阻滞,其中,男 27 例,女 14 例,平均年龄( $5.6 \pm 1.7$ )岁,两组患儿病例资料具有可比性( $P>0.05$ )。

### 1.2 麻醉与手术方法

所有患儿麻醉前禁食 6 h,禁饮 4 h,入室后常规持续监测血压、心率、血氧饱和度。

观察组:面罩持续吸入 8%七氟醚 1-2 min,入睡后建立静脉通道,观察患儿睫毛反射情况,待反射消失后,置入喉罩并固定,七氟醚调整 2%-3%,氧流量 1 L/min,1 人辅助让患儿取左侧卧位,双腿屈曲,颈部不曲,定位骶管并标记,常规消毒铺巾,进行骶管穿刺,骶管穿刺成功后,注入 1%利多卡因 0.5 mL/kg+0.3%罗哌卡因 0.5 mL/kg 的混合液,注射完毕后患儿取平卧位,保持呼吸顺畅。采用传统疝囊高位结扎术进行治疗,下腹部皮肤正中横纹处取手术切口,逐层切开皮肤,显露精索,游离疝囊至疝颈部,8 字缝合后高位结扎,逐层缝合手术切口。术中麻醉维持剂量:2%-3%七氟醚,氧流量 1 L/min,根据麻醉程

度调整七氟醚浓度。手术结束前 10 min 停止吸入七氟醚。

对照组:患儿入室后,从静脉给予阿托品 0.01 mg/kg,氯胺酮 2 mg/kg,面罩吸氧 2 L/min,2 min 后行手术,也采用传统疝囊高位结扎术进行治疗,术中按 1 mg/kg 间断追加,术前 10 min 停止静脉注射氯胺酮。

### 1.3 观察指标

①生命体征监测:观察并记录两组患儿心率(HR),平均动脉压(MAP),血氧饱和度( $SpO_2$ ),分别在切皮前 5 min( $T_0$ )、切皮后 5 min( $T_1$ )、切皮后 10 min( $T_2$ )及切皮后 15 min( $T_3$ )四个时间点记录;②观察并记录两组患儿麻醉诱导时间、苏醒时间和手术麻醉时间;③苏醒期躁动评分(Pediatric anesthesia emergence delirium, PAED)和麻醉诱导期合作量表(Induction Compliance Checklist, ICC);PAED 可以用于评估小儿麻醉术后躁动情况,量表包括 5 项,满分 20 分,>16 分记为苏醒期躁动。ICC 用于评定患儿麻醉诱导时的行为表现,满分 10 分,得分越低说明小儿合作程度越好;④不良反应:观察并记录麻醉期间两组不良反应情况。

### 1.4 统计方法

采用 SPSS 17.0 统计软件分析,数据以  $\bar{x} \pm s$  表示,组内术前与术后相比采用配对 t 检验,组间比较采用两独立样本 t 检验,计数资料采用卡方检验,以  $P<0.05$  为差异有统计学意义。

## 2 结果

### 2.1 HR, MAP, $SpO_2$ 比较

两组切皮后  $T_1$ 、 $T_2$  时 HR、MAP 水平均高于  $T_0$  时的值( $P<0.05$ ),两组切皮后  $T_1$ 、 $T_2$  时组间 HR、MAP 水平相比,无统计学差异( $P>0.05$ )。两组切皮后  $T_3$  时 HR、MAP 水平基本恢复到  $T_0$  时的水平。两组切皮前后 4 个时间点的  $SpO_2$  相比,无统计学差异( $P>0.05$ ),见表 1。

表 1 两组 HR, MAP,  $SpO_2$  指标对比 ( $\bar{x} \pm s$ )

Table 1 Comparison of HR, MAP,  $SpO_2$  between two groups ( $\bar{x} \pm s$ )

| Groups                    | Indexes       | $T_0$           | $T_1$            | $T_2$            | $T_3$           |
|---------------------------|---------------|-----------------|------------------|------------------|-----------------|
| Observation group<br>N=41 | HR (time/min) | 106.7 $\pm$ 5.9 | 127.1 $\pm$ 6.9* | 120.5 $\pm$ 5.7* | 110.8 $\pm$ 3.9 |
|                           | MAP (mm Hg)   | 56.5 $\pm$ 3.7  | 62.8 $\pm$ 4.7*  | 64.1 $\pm$ 6.1*  | 57.8 $\pm$ 7.3  |
|                           | $SpO_2$ (%)   | 97.6 $\pm$ 1.4  | 96.5 $\pm$ 1.3   | 95.8 $\pm$ 1.8   | 97.1 $\pm$ 1.1  |
| Control group<br>N=39     | HR (time/min) | 107.2 $\pm$ 5.5 | 126.4 $\pm$ 6.6* | 119.4 $\pm$ 4.7* | 109.3 $\pm$ 3.9 |
|                           | MAP (mm Hg)   | 56.2 $\pm$ 3.4  | 61.2 $\pm$ 3.6*  | 62.4 $\pm$ 4.1*  | 56.2 $\pm$ 4.2  |
|                           | $SpO_2$ (%)   | 97.4 $\pm$ 2.1  | 97.6 $\pm$ 1.7   | 97.1 $\pm$ 1.5   | 96.7 $\pm$ 1.4  |

Note: compared with  $T_0$ , \* $P<0.05$ .

### 2.2 麻醉诱导及苏醒时间比较

观察组患儿麻醉诱导时间,苏醒时间均明显短于对照组患

儿( $P<0.05$ ),两组术中麻醉持续时间相比,无统计学差异( $P>0.05$ ),均能达到期望麻醉时间,见表 2。

表 2 两组麻醉效果比较 ( $\bar{x} \pm s$ )

Table 2 Comparison of anesthetic effect between two groups ( $\bar{x} \pm s$ )

|                   | n  | Anesthesia induction time(s) | Recovery time (min) | Anesthesia duration time (min) |
|-------------------|----|------------------------------|---------------------|--------------------------------|
| Observation group | 41 | 48.6 $\pm$ 10.2*             | 7.9 $\pm$ 2.5*      | 33.8 $\pm$ 3.4                 |
| Control group     | 39 | 86.2 $\pm$ 9.7               | 18.7 $\pm$ 5.2      | 31.6 $\pm$ 3.5                 |

Note: compared with the control group, \* $P<0.05$ .

### 2.3 PAED 和 ICC 评分比较

( $P < 0.05$ ), 见表 3。

观察组患儿 PAED 评分和 ICC 评分均低于对照组患儿

表 3 两组 PAED 和 ICC 评分对比 ( $\bar{x} \pm s$ )  
Table 3 Comparison of PAED and ICC between two groups ( $\bar{x} \pm s$ )

| Groups            | n  | PAED            | ICC              |
|-------------------|----|-----------------|------------------|
| Observation group | 41 | 15.2 $\pm$ 4.7* | 1.71 $\pm$ 0.52* |
| Control group     | 39 | 18.1 $\pm$ 5.6  | 2.89 $\pm$ 0.74  |

Note: compared with the control group, \* $P < 0.05$ .

### 2.4 不良反应比较

麻醉期间, 观察组患儿 1 例体温升高, 1 例恶心呕吐, 不良反应率 4.9%, 对照组患儿 1 例恶心呕吐, 不良反应率 2.6%, 组间比较无统计学差异( $P > 0.05$ )。

### 3 讨论

腹股沟疝是小儿常见的外科疾病, 多发生在 3 个月以上的小儿, 该疾病几乎不能自愈, 而采用手术治疗的方式可以取得较为满意的治疗结果<sup>[5,6]</sup>。其中, 小儿疝囊高位结扎术是常用手术方式之一, 用于治疗双侧斜疝或隐性疝, 可降低疝复发和腹腔内粘连<sup>[7,8]</sup>。该种手术术前麻醉多采用局部麻醉或半身麻醉, 较少用全身麻醉。2-6 岁的小儿由于其特殊的生理结构, 麻醉既要考虑麻醉有效性又要考虑安全性<sup>[9,10]</sup>。

手术要求麻醉诱导时间短, 麻醉深度可调控, 麻醉苏醒期短。七氟醚是临床使用的吸入性麻醉药物, 对小儿呼吸道刺激小, 无损伤, 再加上七氟醚本身具有一定芳香气味, 小儿普遍易于接受<sup>[11-13]</sup>。七氟醚血气分布系数仅为 0.63, 麻醉药物能迅速在肺泡进行扩散, 麻醉诱导时间短, 麻醉深度可调控。七氟醚本身理化性质稳定, 用药安全性高, 即便小儿吸入高浓度七氟醚, 也没有出现屏气、咳嗽、喉痉挛等不良反应<sup>[14-16]</sup>。除了吸入麻醉方式, 骶管阻滞麻醉也是小儿外科手术常用的麻醉方式, 麻醉效果可靠且易于操作, 成功率高。2-6 岁小儿骶骨没有骨性融合, 小儿骶裂孔大, 容易穿刺, 该种麻醉方法的镇痛和肌松程度均达到疝气手术需要的效果<sup>[17,18]</sup>。

刘瑶等<sup>[19]</sup>研究发现七氟醚复合骶管阻滞进行麻醉对小儿生命体征影响较小。本研究中两组切皮后 T1、T2 时 HR、MAP 水平均高于 T0 时的值 ( $P < 0.05$ ), 两组切皮后 T1、T2 时组间 HR、MAP 水平相比, 无统计学差异( $P > 0.05$ )。两组切皮后 T3 时 HR、MAP 水平基本恢复到 T0 时的水平。两组切皮后四个时间点的 SpO<sub>2</sub> 相比, 无统计学差异( $P > 0.05$ )。七氟醚麻醉对交感神经的兴奋作用较弱, 故七氟醚联合骶管阻滞麻醉对小儿生命体征影响较小。孙红梅等<sup>[20]</sup>研究发现采用骶管阻滞复合七氟醚麻醉 25 例患儿, 平均苏醒时间为(11.0 $\pm$  4.0) min。观察组患儿麻醉诱导时间(48.6 $\pm$  10.2) min, 苏醒时间(7.9 $\pm$  2.5) min, 均短于对照组患儿( $P < 0.05$ ), 也比孙红梅等报道的苏醒时间短。两组术中麻醉持续时间相比, 无统计学差异( $P > 0.05$ ), 均能达到期望麻醉时间, 说明七氟醚联合骶管阻滞麻醉诱导快, 苏醒快, 麻醉效果与静脉注射氯胺酮相当。观察组患儿 PAED 评分和 ICC 评分均低于对照组患儿( $P < 0.05$ ), 静脉注射氯胺酮, 需要扎针, 故小儿配合度较低。此外, 静脉注射氯胺酮, 小儿苏醒期间躁动频

率增加。采用七氟醚复合骶管阻滞, 不仅小儿配合度有所提高, 苏醒期躁动频率也较低, 与七氟醚药物性质、骶管阻滞麻醉的有效性相关。麻醉期间, 观察组患儿 1 例体温升高, 但术后 12 h 内小儿体温自行恢复正常; 共出现 1 例恶心、呕吐, 给予止吐药缓解症状。

综上所述, 七氟醚联合骶管阻滞麻醉对小儿疝气手术具有良好的麻醉效果, 麻醉诱导快, 苏醒快, 小儿配合度高, 术后躁动少, 值得临床推广使用。

### 参考文献 (References)

- [1] Voorbrood CE, Burgmans JP, Clevers GJ, et al. One-stop endoscopic hernia surgery: efficient and satisfactory [J]. *Hernia*, 2015, 19 (3): 395-400
- [2] Mokini Z, Vitale G, Aletti G, et al. Pain control with ultrasound-guided inguinal field block compared with spinal anesthesia after hernia surgery: a randomized trial [J]. *Surgery*, 2015, 157(2): 304-311
- [3] 刘清玉, 艾伟, 刘波, 等. 异氟醚与七氟醚复合麻醉在小儿全身麻醉中的应用[J]. 现代生物医学进展, 2016, 16(2): 329-331  
Liu Qing-yu, Ai Wei, Liu Bo, et al. Application of Isoflurane and Sevoflurane Compound Anesthesia in Children General Anesthesia [J]. *Progress in Modern Biomedicine*, 2016, 16(2): 329-331
- [4] 刘建伟, 蒲晓斌, 张佳君, 等. 瑞芬太尼复合七氟醚在颅内动脉瘤栓塞术中的临床观察[J]. 现代生物医学进展, 2013, 13(33): 6533-6535  
Liu Jian-wei, Pu Xiao-bin, Zhang Jia-jun, et al. The Clinical Observation of Remifentanyl Combined with Sevoflurane in Intracranial Aneurysm Embolization [J]. *Progress in Modern Biomedicine*, 2013, 13(33): 6533-6535
- [5] Hammady A, Abdelgalil W, Rashed E, et al. Upper urinary tract retroperitoneoscopic surgery under epidural anesthesia: Shifting towards outpatient treatment[J]. *Scand J Urol*, 2015, 49(2): 181-184
- [6] Vu MM, Galiano RD, Souza JM, et al. A multi-institutional, propensity-score-matched comparison of post-operative outcomes between general anesthesia and monitored anesthesia care with intravenous sedation in umbilical hernia repair [J]. *Hernia*, 2016, 20 (4): 517-525
- [7] Frawley G, Bell G, Disma N, et al. Predictors of Failure of Awake Regional Anesthesia for Neonatal Hernia Repair: Data from the General Anesthesia Compared to Spinal Anesthesia Study--Comparing Apnea and Neurodevelopmental Outcomes [J]. *Anesthesiology*, 2015, 123(1): 55-65
- [8] Verstraete L, Becaus N, Swannet H, et al. Long term outcome after lichtenstein hernia repair using general, locoregional or local

- anaesthesia[J]. Acta Chir Belg, 2015, 115(2): 136-141
- [9] Özdemir M, Yanlı PY, Tomruk ŞG, et al. Anaesthesia Management of a Patient with Incidentally Diagnosed Diaphragmatic Hernia During Laparoscopic Surgery [J]. Turk J Anaesthesiol Reanim, 2015, 43(1): 50-54
- [10] Milone M, Di Minno MN, Musella M, et al. Outpatient inguinal hernia repair under local anaesthesia: feasibility and efficacy of ultrasound-guided transversus abdominis plane block [J]. Hernia, 2013, 17(6): 749-755
- [11] Nilsson H, Angerås U, Sandblom G, et al. Serious adverse events within 30 days of groin hernia surgery[J]. Hernia, 2016, 20(3): 377-385
- [12] Bourgon AL, Fox JP, Saxe JM, et al. Outcomes and charges associated with outpatient inguinal hernia repair according to method of anesthesia and surgical approach [J]. Am J Surg, 2015, 209(3): 468-472
- [13] Losey-Flores K, Benzar R, Chan JM, et al. Free hernia surgery for the underserved is possible in the United States [J]. Hernia, 2014, 18(2): 305-310
- [14] Tomak Y, Erdivanli B, Sen A, et al. Effect of cooled hyperbaric bupivacaine on unilateral spinal anesthesia success rate and hemodynamic complications in inguinal hernia surgery [J]. J Anesth, 2016, 30(1): 26-30
- [15] Lambert A, Schälte G, Winter J, et al. Spinal anesthesia for inguinal hernia repair in infants: a feasible and safe method even in emergency cases[J]. Pediatr Surg Int, 2014, 30(10): 1069-1073
- [16] Dahlstrand U, Sandblom G, Ljungdahl M, et al. TEP under general anesthesia is superior to Lichtenstein under local anesthesia in terms of pain 6 weeks after surgery: results from a randomized clinical trial [J]. Surg Endosc, 2013, 27(10): 3632-3638
- [17] Suresh S, Schaldenbrand K, Wallis B, et al. Regional anaesthesia to improve pain outcomes in paediatric surgical patients: a qualitative systematic review of randomized controlled trials [J]. Br J Anaesth, 2014, 113(3): 375-390
- [18] Huntington CR, Wormer BA, Cox TC, et al. Local Anesthesia in Open Inguinal Hernia Repair Improves Postoperative Quality of Life Compared to General Anesthesia: A Prospective, International Study [J]. Am Surg, 2015, 81(7): 704-709
- [19] 刘瑶, 贾洪峰. 七氟醚与氯胺酮复合骶管阻滞在小儿疝气手术中应用比较[J]. 航空航天医学杂志, 2011, 22(10): 1156-1157
- Liu Yao, Jia Hong-feng. Depends on Sevoflurane Inhalation and Intramuscular Injection of Ketamine Anesthesia as a Basis for the Inguinal Hernia Sac High Ligation in Children [J]. Journal of Aerospace Medicine, 2011, 22(10): 1156-1157
- [20] 孙红梅, 杨海扣, 拾翠翠. 骶管阻滞复合七氟醚麻醉在小儿腹腔镜单侧疝囊高位结扎术中的应用[J]. 现代中西医结合杂志, 2010, 19(35): 4595-4596
- Sun Hong-mei, Yang Hai-kou, Shi Cui-cui. Application of sacral block combined with sevoflurane anesthesia in children with unilateral laparoscopic high ligation of hernial sac[J]. Modern Journal of Integrated Traditional Chinese and Western Medicine, 2010, 19(35): 4595-4596

## (上接第 1933 页)

- [16] Germain M, Pezzolesi MG, Sandholm N, et al. SORBS1 gene, a new candidate for diabetic nephropathy: results from a multi-stage genome-wide association study in patients with type 1 diabetes [J]. Diabetologia, 2015, 58(3): 543-538
- [17] Kim CS, Park JS, Ahn CW, et al. All-Trans Retinoic Acid Has a Potential Therapeutic Role for Diabetic Nephropathy[J]. Yonsei Med J, 2015, 56(6): 1597-1603
- [18] 段伟喆, 赵湜.  $\alpha$ -硫辛酸联合替米沙坦治疗早期 2 型糖尿病肾脏疾病的临床疗效观察[J]. 临床肾脏病杂志, 2014, 14(7): 415-419
- Duan Wei-zhe, Zhao Shi. Clinical efficacy of  $\alpha$ -lipoic acid combined with telmisartan in treating early type 2 diabetic kidney disease [J]. J Clin Nephrol, 2014, 14(7): 415-419
- [19] Ni WJ, Tang LQ, Wei W. Research progress in signalling pathway in diabetic nephropathy[J]. Diabetes Metab Res Rev, 2015, 31(3): 221-233
- [20] 赵红霞, 董艳双, 蔡友治, 等. 肥胖产妇血脂水平与 DNA 损伤及总抗氧化能力水平的相关性研究 [J]. 医学研究生学报, 2015, 28(2): 157-160
- Zhao Hong-xia, Dong Yan-shuang, Cai You-zhi, et al. Correlation of lipid profile levels with deoxyribonucleic acid damage and total antioxidant capacity levels in obese pregnant women [J]. Journal of Medical Postgraduates, 2015, 28(2): 157-160
- [21] 李玉飞, 鹿玲. 过敏性紫癜患儿血浆髓过氧化物酶、丙二醛、超氧化物歧化酶及总抗氧化能力水平变化[J]. 临床儿科杂志, 2015, 33(4): 357-360
- Li Yu-fei, Lu Ling. Plasma myeloperoxidase, malondialdehyde, superoxide dismutase, and total anti-oxidant capability of Henoch-Schonlein purpura in children [J]. Journal of Clinical Pediatrics, 2015, 33(4): 357-360
- [22] 林杏娟.  $\alpha$ -硫辛酸联合缬沙坦对糖尿病肾病氧化应激的调节作用[J]. 国际医药卫生导报, 2013, 19(8): 1161-1165
- Lin Xing-juan. Effects of  $\alpha$ -lipoic acid and valsartan on the oxidative stress in diabetic nephropathy [J]. International Medicine and Health Guidance News, 2013, 19(8): 1161-1165