

doi: 10.13241/j.cnki.pmb.2018.19.022

老年脓毒症和脓毒性休克患者 30 天死亡率的危险因素分析 *

孙守松¹ 张鑫雨² 陈 静³ 陈小凤¹ 季娟娟¹ 孙 勇¹ 尹琮玉¹ 王 刚^{1Δ}

(1 解放军第四五四医院急诊科 江苏 南京 210000; 2 空军军医大学基础医学院学员四大队 陕西 西安 710032;

3 泰州职业技术学院护理系 江苏 泰州 225300)

摘要 目的:探讨就诊于急诊室诊断为脓毒症或者脓毒性休克的患者 30 天内死亡率与在急诊室初次获得的参数的关系,并探讨死亡率的危险因素。方法:选取 2014 年 6 月到 2017 年 6 月就诊于我院急诊室且诊断为脓毒症或者脓毒性休克并有完整随访资料大于 65 岁的老年患者 145 例,将 30 天内存活的患者分为 A 组,将 30 天内死亡的患者分为 B 组,比较两组之间检验指标及生命体征的差异。根据脓毒症序贯器官衰竭评估快速评分(qSOFA)将 qSOFA<2 分的定义为 a 组,qSOFA≥2 分的定义为 b 组,比较两组的死亡率。并采用二项 Logistic 回归分析探讨 30 天死亡率的独立危险因素。结果:145 例患者中,44.8%(n=65)的患者在 30 天内死亡,33.1%(n=48)的患者需要无创或者侵入性机械通气。A 组与 B 组之间舒张压(P=0.003)、收缩压(P=0.002)、格拉斯哥昏迷量表评分(GCS)(P<0.001)、血尿素氮(P<0.001)及 qSOFA(P<0.001)比较均具有统计学差异。a 组死亡率(35.7%)显著低于 b 组(53.3%)(P=0.033)。qSOFA 是 30 天内死亡率的独立危险因素(OR=2.871, P=0.004)。结论:qSOFA 是老年脓毒症及脓毒性休克患者 30 天内死亡的独立危险因素。

关键词:急诊;老年;脓毒症;30 天内死亡率;危险因素

中图分类号:R459.7;R631.2 文献标识码:A 文章编号:1673-6273(2018)19-3698-04

Analysis of the Risk Factors of 30-day Mortality in Elderly Patients with Sepsis and Septic Shock*

SUN Shou-song¹, ZHANG Xin-yu², CHEN Jing³, CHEN Xiao-feng¹, JI Juan-juan¹, SUN Yong¹, YIN Qiong-yu¹, WANG Gang^{1Δ}

(1 Department of Emergency, 454th Hospital of PLA, Nanjing, Jiangsu, 210000, China; 2 Fourth Cadet Brigade, Air Force Medical University, Xi'an, Shaanxi, 710032, China; 3 Department of nursing, Taizhou Polytechnic College, Taizhou, Jiangsu, 225300, China)

ABSTRACT Objective: To investigate the relationship between 30-day mortality of patients diagnosed with sepsis or septic shock in the emergency room and the initial parameters obtained in the emergency room, and explore the risk factors of mortality. **Methods:** 145 cases of patients older than 65 years old who were diagnosed as sepsis or septic shock and had complete follow-up data were collected from June 2014 to June 2017 in the emergency room of our hospital were selected. The patients who survived within 30 days were divided into group A, and the patients who died within 30 days were divided into group B. The differences in laboratory inspection parameters and vital signs groups were compared between the two. According to the quick Sepsis-related Organ Failure Assessment (qSOFA), patients with qSOFA<2 points were defined as group a, and patients with qSOFA≥2 points were defined as group b. The mortality was compared between the two groups. Binary logistic regression analysis was used to evaluate independent risk factors for 30-day mortality. **Results:** Of the 145 cases of patients, 44.8% (n=65) of patients died within 30 days, and 33.1% (n=48) of patients required non-invasive or invasive mechanical ventilation. There were statistically significant differences in the diastolic blood pressure (P=0.003), systolic blood pressure (P=0.002), Glasgow Coma Scale (GCS) (P<0.001), blood urea nitrogen (P<0.001), and qSOFA (P<0.001) between group A and group B. The mortality rate of group a (35.7%) was significantly lower than that of group b (53.3%)(P=0.033). qSOFA was an independent risk factor for the mortality within 30 days (OR=2.871, P=0.004). **Conclusions:** qSOFA is an independent risk factor for the death within 30 days of sepsis and septic shock patients.

Key words: Emergency; Elderly; Sepsis; Mortality within 30 days; Risk factors

Chinese Library Classification (CLC): R459.7; R631.2 **Document code:** A

Article ID: 1673-6273(2018)19-3698-04

前言

近年来,就诊于急诊室大于 65 岁的老年人逐年增多,在急诊室对于脓毒症老年患者的管理引起学者的关注^[1]。有学者认

* 基金项目:南京军区科研基金面上课题(12MA043)

作者简介:孙守松(1982-),本科,从事急诊外科研究,E-mail:42462878@qq.com

Δ 通讯作者:王刚(1978-),本科,从事急诊外科研究,E-mail:305524361@qq.com,电话:02580865081

(收稿日期:2018-03-17 接受日期:2018-04-20)

为升高的乳酸有助于老年患者脓毒症的诊断及确诊^[4]。基于2016年脓毒症及脓毒性休克的定义,有学者提出脓毒症序贯器官衰竭评估(Sepsis-related Organ Failure Assessment, SOFA)评分系统^[5]。后来又形成了一个在急诊室更简单实用的系统-脓毒症序贯器官衰竭评估快速评分(quick SOFA, qSOFA)。qSOFA结合了感染的特征及另外三个标准:收缩压 ≤ 100 mmHg、格拉斯哥昏迷量表评分(Glasgow Coma Scale, GCS) < 15 和呼吸频率 ≥ 22 次/分,每个标准为1分,如果总分 ≥ 2 分常常提示感染已危及生命。有研究表明qSOFA在没有实验室检查的条件下有助于脓毒症的诊断^[6]。qSOFA ≥ 2 分且合并有需要缩血管药物维持平均动脉压 > 75 mmHg和乳酸 > 2 mmol/L称为脓毒性休克。

除了乳酸水平,有些生物指标比如降钙素原和C反应蛋白也用于诊断和预测脓毒症^[6]。但这些实验室检验指标比全血细胞计数昂贵。血小板与淋巴细胞比(platelet-lymphocyte ratio, PLR)是一个可以从血常规中获得不需要额外花费的指标,有研究报道PLR是炎症反应的象征^[7]。最近的研究表明PLR在各种条件下均升高,包括急性肠系膜缺血、慢性阻塞性肺病急性加重、类风湿性关节炎和肾细胞癌^[8,9]。因此,本研究主要探讨了qSOFA、PLR、乳酸、其它检验指标和生命体征是否是脓毒症和脓毒性休克患者30天内死亡的危险因素,以期老年脓毒症和脓毒性休克患者的诊治提供参考。

1 资料与方法

1.1 一般资料

选取2014年6月到2017年6月就诊于我院急诊室且初次诊断为脓毒症或者脓毒性休克大于65岁的老年患者。通过医院门诊系统检索患者的基本资料、初次实验室检验指标及生命体征。通过电话回访患者在我院急诊确诊后的30天内的生存状态。将30天内存活的患者分为A组(80例),将30天内死亡的患者分为B组(65例)。

1.2 观察指标

患者的基本资料:年龄、性别及合并症。入急诊室时的收缩压、舒张压、GCS、qSOFA、呼吸频率、体温、心率、氧饱和度、血钠、血钾、血尿素氮、血肌酐、血红蛋白、血乳酸、白细胞、血小板、中性粒细胞、淋巴细胞和PLR。比较上述指标在A组与B组之间的差异。此外,收集患者在急诊室接受的治疗。

1.3 统计学分析

通过SPSS20.0分析数据,计量资料以均值 $\pm$ 标准差表示,计数资料以百分数表示。分别利用t检验及卡方检验,采用二项Logistic回归分析探讨死亡率的独立危险因素。以 $P < 0.05$ 为差异有统计学意义。

2 结果

2.1 一般情况

从2014年6月到2017年6月,共有145例患者纳入研究,年龄为 82.0 ± 10.5 ,其中女性占54.5%(79例),男性占45.5%(66例)。44.8%(65例)的患者在入急诊室30天内死亡。145例患者中,最常见的感染原因为肺部感染(69.0%)和泌尿系统感染(46.2%),最常见的合并症是高血压(62.1%)、冠心病(29.0%)

和糖尿病(27.6%)。在急诊室接受最常见的治疗为吸氧(59.3%)、抗菌药使用(55.2%)和支气管舒张剂的使用(53.1%),见表1。

2.2 A组与B组之间各项指标的差异

表1 在急诊室接受的治疗

Table 1 Treatment started in the emergency room

Treatment	%, n
Oxygen absorption	59.3(86)
Antibacterial drugs	55.2(80)
Bronchodilator	53.1(77)
Intravenous hydration	40.7(59)
Paracetamol	33.8(49)
Mechanical Ventilation	33.1(48)
Furosemide	28.3(41)
Methylprednisolone	16.6(24)
Low molecular weight heparin	11.7(17)
Dopamine	11.0(16)
Calcium antagonists	8.3(12)
Cardiopulmonary resuscitation	5.5(8)
Nitroglycerin	4.8(7)

A组与B组之间各个参数的差异见表2。A组的收缩压($P=0.002$)、舒张压($P=0.003$)和GCS($P < 0.001$)均高于B组,血尿素氮($P < 0.001$)及qSOFA($P < 0.001$)均低于B组,差异具有统计学意义。两组之间的血肌酐、PLR和其他参数比较均没有差异。

2.3 不同qSOFA评分患者的死亡率

将qSOFA < 2 分的患者分为a组,将qSOFA ≥ 2 分的患者分为b组,b组死亡率(53.3%)显著高于a组(35.7%)($P=0.033$),见表3。

2.4 患者生存状态的多因素分析

以是否死亡为因变量,以表1中A组与B组有差异的参数(收缩压、舒张压、GCS、血尿素氮和qSOFA)为自变量,进行二项Logistic回归分析,结果显示qSOFA为30天内死亡的独立危险因素($OR=2.871$, $P=0.004$),见表4。

3 讨论

脓毒症和脓毒性休克在全世界范围内均是死亡率很高且治疗费用很高的疾病^[1]。老年脓毒症患者的临床特征年轻患者不同^[10]。本研究主要探讨了PLR、乳酸及其他指标能否用于诊断脓毒症,及患者30天死亡率的独立危险因素。结果显示存活组与死亡组的PLR和乳酸水平比较均没有显著差异,但两组患者的qSOFA不同,不同qSOFA的患者死亡率不同,且qSOFA是30天内死亡的独立危险因素。

Safari的研究表明在急诊室确诊的脓毒症或者脓毒性休克的患者30天死亡率为51.43%,这些患者的年龄为 68.36 ± 18.62 ^[11]。Chen通过评估qSOFA、CRB-65和CRB在肺炎中的预测性能,发现28天的死亡率为33%^[12]。本研究中,患者30

天死亡率为 44.8 %，和上述结果相似,33.1 %的患者需要机械通气^[14],Dettmer 同样报道了在急诊室中需要机械通气的患者比例为 39.9 %^[15]。

Bhat 报道了诊断为严重脓毒症的患者中有 41.5 %需要机械通

表 2 A组与 B 组参数比较

Table 2 Comparison of the parameters between group A and group B

Parameters	Group A(n=80)	Group B(n=65)	P value
Diastolic pressure(mmHg)	71.36± 22.41	61.34± 20.65	0.003
Systolic pressure(mmHg)	130.21± 36.12	110.25± 38.42	0.002
GCS(score)	14.3± 2.1	12.7± 1.8	<0.001
Blood urea nitrogen(mg/dl)	26.2± 7.6	47.6± 10.1	<0.001
qSOFA(score)	1.73± 0.32	2.21± 0.41	<0.001
Age(year)	81.9± 7.8	82.1± 7.9	0.879
Gender(Female,%)	42, 52.5	37, 56.9	0.595
Breathing frequency(/min)	22.1± 7.1	23.7± 7.6	0.193
Body temperature(℃)	36.9± 1.0	37.1± 1.1	0.254
Heart rate(/min)	99.8± 23.5	103.7± 24.1	0.328
SpO ₂ (%)	85.7± 12.7	83.8± 11.9	0.358
Creatinine(mg/dl)	140.7± 21.7	145.9± 23.7	0.171
Blood sodium(mmol/L)	135.3± 10.7	136.7± 11.1	0.442
Blood potassium(mmol/L)	4.31± 0.78	4.41± 0.73	0.431
Hemoglobin (g/dL)	12.79± 2.56	12.34± 2.76	0.322
White blood cells (thousands/ μ L)	12.37± 7.12	13.64± 7.03	0.285
Platelets (thousands/ μ L)	212.00± 97.65	215.81± 98.67	0.816
Neutrophils (thousands/ μ L)	10.17± 6.47	11.87± 7.12	0.135
Lymphocytes (thousands/ μ L)	0.94± 0.51	0.97± 0.54	0.732
Blood lactate (mmol/L)	1.97± 0.86	2.26± 0.98	0.060
PLR	204.7± 74.8	189.6± 63.4	0.198

表 3 a 组与 b 组死亡率的比较

Table 3 Comparison of the mortality rate between group a and group b

Living status	Group a (n, %)	Group b (n, %)	P value
Death	25, 35.7	40, 53.3	0.033
Survival	45, 64.3	35, 46.7	

表 4 Logistic 回归分析结果

Table 4 Logistic regression analysis results

Parameters	B value	SE	Wals	P value	Exp(B)
qSOFA	1.055	0.365	8.326	0.004	2.871
Systolic pressure	0.012	0.018	0.441	0.507	1.012
Diastolic pressure	0.150	0.180	0.691	0.406	1.162
GCS	0.286	0.615	0.215	0.643	1.331
Blood urea nitrogen	1.368	0.886	2.388	0.122	3.927

Note: P-value is the result of comparative analysis between group A and group B.

Valencia 也报道了 >65 岁且诊断为脓毒症的患者在急诊室中,存活的患者平均动脉压高于死亡的患者^[10]。脓毒症或脓毒性休克的患者由于炎性因子的大量释放,引起血管通透性的增加及血管对缩血管的敏感性降低,引起血压下降,血压下降

后减少了心、肾、脑及外周血管的氧供,造成组织缺氧。故存活的患者血压往往较死亡者更高。本研究中,存活组与死亡组的收缩压与舒张压具有显著差异,且两组患者的GCS评分和血尿素氮水平不同。Nowak研究了在急诊室诊断为脓毒症的患者,发现低死亡率和高死亡率的患者血尿素氮水平没有统计学差异,我们的结果与上述结果不同有可能是因为两个研究纳入的人群不同,Nowak研究的人群为<65岁的患者^[16]。在另外一个和我们设计相似的研究中,Uluöz发现死亡患者的GCS(10.36 ± 4.9)评分显著低于存活的病人(13.6 ± 2.5)^[17],和我们的研究结果一致。

肺部感染和泌尿系统感染是脓毒症最常见的病因^[18,19]。最常见的三个合并症是高血压(62.1%)、冠心病(29.0%)和糖尿病(27.6%)。Logoglu发现急诊室中>65岁的患者最常见合并症分别是高血压(40.5%)、冠心病(26.6%)和糖尿病(22.4%)^[20],和我们的结果相似。抗菌药是在急诊室用于治疗最常用的药物(55.2%),与我们的结果一致^[21]。本研究中,死亡组qSOFA评分(2.21 ± 0.41)显著高于存活组(1.73 ± 0.32),qSOFA ≥ 2 分的患者死亡率(53.3%)显著高于qSOFA<2分(35.7%)的患者,qSOFA与死亡率存在显著的正相关。此外,qSOFA评分是死亡率的独立危险因素。Wang对477例急诊患者的qSOFA评分进行分析,发现存活组与死亡组的qSOFA评分显著不同^[22]。Singer的研究发现qSOFA评分与住院率、住院时间和住院期间死亡率呈现显著正相关^[23]。我们认为感染越重的患者,组织缺氧和酸中毒就越重,更易出现休克、血压下降和呼吸急促,故qSOFA评分往往就更高。Zencir发现在致死性感染心内膜炎的患者中PLR升高^[24]。然而Duman发现在脓毒症、脓毒性休克和严重的脓毒症之间PLR没有差异($P=0.737$),在存活和死亡的病人中也没有发现PLR有任何差异($P=0.336$)^[25]。虽然PLR被认为是一个经济且易获得的感染指标,但是我们和Duman的研究均没有发现死亡和存活的患者PLR有差异。在未来的研究中,仍需要更多研究PLR与感染及愈后的关系。

此外,我们还发现存活组与死亡组之间的初次检验的乳酸值没有统计学差异。Varis发现入院时乳酸>2 mmol/L的患者90天死亡率大于乳酸<2 mmol/L的患者,但是作者发现入院时的乳酸值不是死亡率的独立危险因素^[26]。结合我们的研究结果,我们认为在病程中多次测量乳酸可能比入院时单次的乳酸测量更能预测患者的愈后。

总之,本研究结果表明qSOFA是老年脓毒症及脓毒性休克患者30天内死亡的独立危险因素,应引起临床医生的足够重视。

参考文献(References)

- [1] 董丽华,吕娟,丁黎莉,等.脓毒症免疫治疗的研究进展[J].中华危重病急救医学,2017,29(2): 184-187
Dong Li-hua, Lv Juan, Ding Li-li, et al. Advances in immunotherapeutic research of sepsis[J]. Chinese Critical Care Medicine, 2017, 29(2): 184-187
- [2] Gaieski DF, Edwards JM, Kallan MJ, et al. Benchmarking the incidence and mortality of severe sepsis in the United States[J]. Crit Care Med, 2013, 41(5): 1167-1174
- [3] Özel BA, Ekinçi EM, Kayipmaz AE, et al. Analysis of the use of resources and features of presentations and the trends in geriatric patients presenting to the emergency department: 2011-2015 [J]. Turk J Geriatr, 2016, 19: 154-161
- [4] Mikkelsen ME, Miltiades AN, Gaieski DF, et al. Serum lactate is associated with mortality in severe sepsis independent of organ failure and shock[J]. Crit Care Med, 2009, 37(5): 1670-1677
- [5] Seymour CW, Liu VX, Iwashyna TJ, et al. Assessment of Clinical Criteria for Sepsis: For the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3)[J]. JAMA, 2016, 315: 801-810
- [6] Green JP, Berger T, Garg N, et al. Serum lactate is a better predictor of short-term mortality when stratified by C-reactive protein in adult emergency department patients hospitalized for a suspected infection [J]. Ann Emerg Med, 2011, 57(3): 291-295
- [7] Kutlucan L, Kutlucan A, Basaran B, et al. The predictive effect of initial complete blood count of intensive care unit patients on mortality, length of hospitalization, and nosocomial infections [J]. Eur Rev Med Pharmacol Sci, 2016, 20(8): 1467-1473
- [8] Toptas M, Akkoc j, Savas Y, et al. Novel hematologic inflammatory parameters to predict acute mesenteric ischemia [J]. Blood Coagul Fibrinolysis, 2016, 27(2): 127-130
- [9] Kilic E, Rezvani A, Toprak AE, et al. Evaluation of Neutrophil to Lymphocyte and Platelet to Lymphocyte Ratios in Rheumatoid Arthritis[J]. Dicle Med J, 2016, 43(2): 241-247
- [10] Valencia AM, Vallejo CE, Alvarez AL, et al. Attenuation of the physiological response to infection on adults over 65 years old admitted to the emergency room (ER)[J]. Aging Clin Exp Res, 2016, 29(5): 847-856
- [11] Safari S, Shojaei M, Rahmati F, et al. Accuracy of SOFA score in prediction of 30-day outcome of critically ill patients [J]. Turk J Emerg Med, 2016, 16(4): 146-150
- [12] Chen YX, Wang JY, Guo SB. Use of CRB-65 and quick Sepsis-related Organ Failure Assessment to predict site of care and mortality in pneumonia patients in the emergency department: a retrospective study[J]. Crit Care, 2016, 20(1): 167
- [13] Sarkari NN, Perman SM, Ginde AA. Impact of early do-not-attempt-resuscitation orders on procedures and outcomes of severe sepsis[J]. J Crit Care, 2016, 36: 134-139
- [14] Bhat SR, Swenson KE, Francis MW, et al. Lactate Clearance Predicts Survival Among Patients in the Emergency Department with Severe Sepsis[J]. West J Emerg Med, 2015, 16(7): 1118-1126
- [15] Dettmer MR, Mohr NM, Fuller BM. Sepsis-associated pulmonary complications in emergency department patients monitored with serial lactate: An observational cohort study [J]. J Crit Care, 2015, 30(6): 1163-1168
- [16] Nowak RM, Reed BP, Nanayakkara P, et al. Presenting hemodynamic phenotypes in ED patients with confirmed sepsis [J]. Am J Emerg Med, 2016, 34(12): 2291-2297
- [17] Uluöz HO, Sebe A, Ay MO, et al. The Relationship Between Inflammatory Reagents and Mortality in Patients Over the Age of 55 Hospitalised in the Internal Medicine Intensive Care Unit from the Emergency Service[J]. JAEM, 2013, 12(1): 13-18
- [18] Carpio R, Zapata J, Spanuth E, et al. Utility of presepsin (sCD14-ST) as a diagnostic and prognostic marker of sepsis in the emergency department[J]. Clin Chim Acta, 2015, 450(1): 169-175 (下转第 3741 页)

- [17] Shin S, Kim S. Dental treatment in patients with severe gag reflex using propofol-remifentanyl intravenous sedation[J]. *J Dent Anesth Pain Med*, 2017, 17(1): 65-69
- [18] Shin DW, Cho JY, Han YS, et al. Risk factor analysis of additional administration of sedative agent and patient dissatisfaction in intravenous conscious sedation using midazolam for third molar extraction [J]. *J Korean Assoc Oral Maxillofac Surg*, 2017, 43(4): 229-238
- [19] Wang J, Han Z, Zhou H, et al. Effective Loading Dose of Dexmedetomidine to Induce Adequate Sedation in Parturients Undergoing Cesarean Section Under Spinal Anaesthesia[J]. *Turk J Anaesthesiol Reanim*, 2017, 45(5): 260-263
- [20] G K, Kakade PK, Singh M, et al. Assessment of Alteration in Capnometry Monitoring during Intravenous Sedation with Midazolam for Oral Surgical Procedures [J]. *J Contemp Dent Pract*, 2017, 18(11): 1025-1028
- [21] Mishra N, Birmiwai KG, Pani N, et al. Sedation in oral and maxillofacial day care surgery: A comparative study between intravenous dexmedetomidine and midazolam[J]. *Natl J Maxillofac Surg*, 2016, 7(2): 178-185
- [22] 许涛,杨涛,曹好好,等.咪达唑仑或丙泊酚联合右美托咪啶对颅脑创伤患者镇静作用的比较 [J]. *中国医院药学杂志*, 2015, 35(4): 326-329
- Xu Tao, Yang Tao, Cao Hao-hao, et al. Comparative analysis of sedative effects mediated by dexmedetomidine in combination with propofol or midazolam in traumatic brain injury patients [J]. *Chinese Journal of Hospital Pharmacy*, 2015, 35(4): 326-329
- [23] Saiso K, Adnonla P, Munsil J, et al. Complications associated with intravenous midazolam and fentanyl sedation in patients undergoing minor oral surgery[J]. *J Dent Anesth Pain Med*, 2017, 17(3): 199-204
- [24] Lovett P, Gómez V, Hodge DO, et al. Propofol Versus Midazolam/Fentanyl Sedation for Colonoscopy in the Elderly Patient Population[J]. *J Perianesth Nurs*, 2017, 32(3): 210-214
- [25] 郝江,罗积慎,翁奇,等.右美托咪定与异丙酚对颅脑创伤患者β-内啡肽水平的影响及镇静疗效对比研究 [J]. *中华危重病急救医学*, 2013, 25(6): 373-376
- Hao Jiang, Luo Ji-shen, Weng Qi, et al. Effects of dexmedetomidine on sedation and β-endorphin in traumatic brain injury: a comparative study with propofol [J]. *Chinese Critical Care Medicine*, 2013, 25(6): 373-376
- [26] 王广,罗超,彭彤,等.脑出血与蛛网膜下腔出血后β-内啡肽和促性腺激素释放激素水平变化研究 [J]. *中国全科医学*, 2017, 20(6): 673-677
- Wang Guang, Luo Chao, Peng Xing, et al. Changes of β-endorphin and Gonadotropin Releasing Hormone after Cerebral Hemorrhage and Subarachnoid Hemorrhage[J]. *Chinese General Practice*, 2017, 20(6): 673-677
- [27] Feng SD, Jiang Y, Lin ZH, et al. Diagnostic value of brain natriuretic peptide and β-endorphin plasma concentration changes in patients with acute left heart failure and atrial fibrillation [J]. *Medicine (Baltimore)*, 2017, 96(34): e7526
- [28] Misra UK, Kalita J, Tripathi G, et al. Role of β endorphin in pain relief following high rate repetitive transcranial magnetic stimulation in migraine[J]. *Brain Stimul*, 2017, 10(3): 618-623
- [29] 黄珩,李萌,何艺,等.丙泊酚联合右美托咪啶对肺癌根治术患者镇痛及应激反应的影响 [J]. *现代生物医学进展*, 2017, 17(13): 2495-2497, 2520
- Huang Heng, Li Meng, He Yi, et al. Analgesic Effects of Propofol Combined with Dexmedetomidine on the Patients Undergoing Radical Resection of Lung Cancer and Its Effect on the Stress Response[J]. *Progress in Modern Biomedicine*, 2017, 17(13): 2495-2497, 2520
- [30] Guldenmund P, Vanhaudenhuyse A, Sanders RD, et al. Brain functional connectivity differentiates dexmedetomidine from propofol and natural sleep[J]. *Br J Anaesth*, 2017, 119(4): 674-684

(上接第 3701 页)

- [19] Mayr FB, Yende S, Angus DC. Epidemiology of severe sepsis[J]. *Virulence*, 2014, 5(1): 4-11
- [20] A L, C A, A K, et al. Analysis of non-traumatic elderly patient presentations to the emergency department[J]. *Turk J Emerg Med*, 2013, 13: 171-179
- [21] Kuan WS, Ibrahim I, Leong BS, et al. Emergency Department Management of Sepsis Patients: A Randomized, Goal-Oriented, Noninvasive Sepsis Trial[J]. *Ann Emerg Med*, 2015, 67(3): 367-368
- [22] Wang JY, Chen YX, Guo SB, et al. Predictive performance of quick Sepsis-related Organ Failure Assessment for mortality and intensive care unit admission in patients with infection at the ED[J]. *Am J Emerg Med*, 2016, 34(9): 1788-1793
- [23] Singer AJ, Ng J, Jr TH, et al. Quick SOFA Scores Predict Mortality in Adult Emergency Department Patients With and Without Suspected Infection[J]. *Ann Emerg Med*, 2017, 68(4): S14-S5
- [24] Zencir C, Akpek M, Senol S, et al. Association between hematologic parameters and in-hospital mortality in patients with infective endocarditis[J]. *Kaohsiung J Med Sci*, 2015, 31(12): 632-638
- [25] Afyon M. Platelet-to-lymphocyte ratio as a cost-effective and easily measurable marker[J]. *Cukurova Medizinhist J*, 2016, 41: 610-611
- [26] Varis E, Pettila V, Poukkanen M, et al. Evolution of Blood Lactate and 90-Day Mortality in Septic Shock. A Post Hoc Analysis of the FINNAKI Study[J]. *Shock*, 2017, 47(5): 574-581